

# COMPACT SPACES, ELEMENTARY SUBMODELS, AND THE COUNTABLE CHAIN CONDITION

LÚCIA R. JUNQUEIRA, PAUL LARSON, AND FRANKLIN D. TALL

September 21, 2004

*Dedicated to Jim Baumgartner on the occasion of his 60th birthday*

ABSTRACT. Given a space  $\langle X, \mathcal{J} \rangle$  in an elementary submodel  $M$  of  $H(\theta)$ , define  $X_M$  to be  $X \cap M$  with the topology generated by  $\{U \cap M : U \in \mathcal{J} \cap M\}$ . It is established, using anti-large-cardinals assumptions, that if  $X_M$  is compact and its regular open algebra is isomorphic to that of a continuous image of some power of the two-point discrete space, then  $X = X_M$ . Assuming  $CH+SCH$  (the Singular Cardinals Hypothesis) in addition, the result holds for any compact  $X_M$  satisfying the countable chain condition.

## 1. INTRODUCTION

This paper continues the line of research of [10], [11], [6], [8] and [12], in which the question of which topological spaces are determined by their compact reflections in elementary submodels is investigated. A minor technical obstacle results from the fact that we cannot take elementary submodels of the entire universe, but we want our models to be elementary in structures much larger than the spaces we are considering. So, we adopt the following convention: whenever  $X$  is a topological space, the elementary submodels we consider have  $X$  as an element and are elementary in  $H(\theta)$  for some regular cardinal  $\theta$  of cardinality greater than all finite iterations of the power-set function starting with  $X$ .

Given a space  $\langle X, \mathcal{T} \rangle$  in an elementary submodel  $M$  of  $H(\theta)$ , we define  $X_M$  to be  $X \cap M$  with the topology generated by  $\{U \cap M : U \in \mathcal{T} \cap M\}$  [5]. If  $X_M$  is compact  $T_2$  (in fact, we shall assume all spaces are  $T_2$ ), this constrains  $X$  to the point that simple additional topological hypotheses on  $X_M$  ensure that  $X_M = X$  [6]. When powers of the two-point discrete space  $D$  are considered, the situation is more complicated: roughly, for  $\kappa$  below very large cardinals,  $X_M$  homeomorphic to  $D^\kappa$  implies  $X_M = X$ , but this is not the case above such large cardinals [8], [11], [6]. This was generalized to continuous images of powers of  $D$  in [12]. In Section 5, we generalize the positive results to compact spaces co-absolute with such spaces. Yet a further generalization is to compact spaces satisfying the countable chain condition. However, for this we need to assume  $CH + SCH$  (where  $SCH$  stands for the Singular Cardinals Hypothesis).

---

*Key words and phrases.* compact, countable chain condition, reflection, elementary submodel, co-absolute with dyadic compact space, squashable.

MSC 2000: Primary 54O30, 54A35, 03E35, 03E55, 03E75; Secondary 54G12, 03G05.

The first author acknowledges support from FAPESP.

The second author acknowledges support from FAPESP (Grant # 02/11551-3) and the Centre de Recerca Matemàtica of the Institut d'Estudis Catalàn.

The third author acknowledges support from NSERC grant A-7354.

In [6] we also considered the contrasting situation of when  $X$  compact implies  $X_M$  compact. This turned out to be related to whether  $X$  is *scattered*, i.e. each subspace has an isolated point. Generalizations of scattering play a key role in the two new theorems mentioned above; as well, we explore in general the relationships between various forms of scattering and the question of “squashing” a compact space  $X$  to a compact  $X_M$ . Formally,

**Definition 1.1.** [8] A compact space  $X$  is *squashable* if for some elementary submodel  $M$  containing  $X$ ,  $X_M$  is compact but not equal to  $X$ .

Kunen [8] noted that squashability does not depend on  $\theta$ . He also showed:

**Lemma 1.2.** *If  $D^\lambda$  is squashable,  $\lambda$  is greater than the first 1-extendible cardinal.*

1-extendible cardinals are reasonably large; in particular, if  $\kappa$  is 1-extendible,  $\kappa$  is the  $\kappa$ th measurable cardinal. For the definition and more on such cardinals, see [7] or [8].

In previous papers, [10], [11], [6], [12], we have used the anti-large-cardinal assumption “ $0^\#$  does not exist” or, rather, its consequence that “ $|M| \geq \kappa$  implies  $M \supseteq \kappa$ ” to limit the types of elementary submodels that can exist. Here we introduce a weaker assumption that will serve our purposes.

**Definition 1.3.** (B): if  $\theta$  is a regular cardinal,  $M$  is an elementary submodel of  $H(\theta)$ , and  $\gamma$  is a cardinal in  $M$  such that  $2^\gamma \in M$ , then  $|M \cap \gamma| \subseteq M$ .

Note that since  $M$  may not satisfy the power set axiom, the condition on  $\gamma$  in the statement of (B) is not vacuous.

**Theorem 1.4.** *Axiom (B) implies no  $D^\kappa$  is squashable.*

*Proof.* We show the contrapositive. Suppose that  $\kappa$  is a cardinal and  $M$  is an elementary submodel of some  $H(\theta)$  such that  $\kappa$  and  $2^\kappa$  are in  $M$  and  $(D^\kappa)_M$  is compact but not equal to  $D^\kappa$ . Then  $\kappa$  is not included in  $M$ , since  $\kappa \subseteq M$  and  $(D^\kappa)_M$  compact imply  $(D^\kappa)_M = D^\kappa$  [11], [6]. Thus we can fix  $\alpha < \kappa$ , the least ordinal not in  $M$ . By our convention, since  $D^\kappa \in M$ ,  $2^{2^\kappa}$  is also in  $M$ . Since  $(D^\kappa)_M$  is compact,  $|2^\kappa \cap M| = 2^{|\kappa \cap M|}$ .  $|\kappa \cap M| \geq |\alpha|$ , so  $2^{|\kappa \cap M|} > \alpha$ . But (B) would imply  $|2^\kappa \cap M| \subseteq M$ , so  $\alpha \in M$ , contradiction.  $\square$

For any elementary submodel  $M$  of any  $H(\theta)$ , we define  $o_M$  to be the least cardinal  $\kappa$  such that  $\kappa^+ \not\subseteq M$  (note that even if  $o_M$  is a limit cardinal,  $M$  must include it). Equivalently,  $o_M$  is the cardinality of the least ordinal not in  $M$ . If  $o_M \neq \theta$  (which must be the case if  $M$  is part of a counterexample to (B)) one of the following must hold:

- $o_M \in M$ , in which case  $o_M^+ \cap M = \eta$ , where  $\eta$  is the least ordinal not in  $M$  (since for every ordinal  $\alpha$  in  $M$  of cardinality  $o_M$  there is a bijection between  $o_M$  and  $\alpha$  in  $M$ );
- $o_M \notin M$ , in which case  $o_M$  is a limit cardinal, and the least ordinal in  $M$  greater than  $o_M$  is a cardinal (as  $M$  is closed under the function  $\alpha \mapsto |\alpha|$ ).

The following reformulation of (B) in terms of  $o_M$  is immediate.

**Theorem 1.5.** *Axiom (B) is equivalent to the assertion that if  $\theta$  is a regular cardinal and  $M$  is an elementary submodel of  $H(\theta)$ , then  $|M \cap \gamma| = o_M$  for every  $\gamma \geq o_M$  in  $M$  such that  $2^\gamma \in M$ .*

Axiom (B) is essentially the principle  $|M| = o_M$  weakened so that, while it suffices for all of our applications, its failure implies a certain form of Chang’s Conjecture (it is for this that

we require  $2^\gamma \in M$ ). Recall that for cardinals  $\lambda > \kappa$  and  $\delta > \eta$ , the expression  $\langle \lambda, \kappa \rangle \rightarrow \langle \delta, \eta \rangle$  says that the set of  $Z \subseteq \lambda$  of cardinality  $\delta$  with  $|Z \cap \kappa| = \eta$  is stationary in  $\mathcal{P}(\lambda)$  (equivalently, that for every function  $F$  from the finite subsets of  $\lambda$  to  $\lambda$  there exists a  $Z \subseteq \lambda$  of cardinality  $\delta$  closed under  $F$  with  $|Z \cap \kappa| = \eta$ ). Another equivalent form of  $\langle \lambda, \kappa \rangle \rightarrow \langle \delta, \eta \rangle$  is the following: there is an elementary substructure  $M$  of  $H(\lambda^+)$  of cardinality  $\delta$  with  $|M \cap \kappa| = \eta$ . *Chang's Conjecture* is the statement  $\langle \omega_2, \omega_1 \rangle \rightarrow \langle \omega_1, \omega \rangle$ .

Supposing that (B) fails, fix the smallest  $\theta$  for which there exists a counterexample, and let  $\zeta$  be the least cardinal for which there exists an  $M \prec H(\theta)$  witnessing the failure of (B) with  $o_M = \zeta$ . Let  $\gamma$  be the least ordinal such that there exists an  $M \prec H(\theta)$  with

- $\gamma, 2^\gamma \in M$ ,
- $\gamma \geq \zeta$ ,
- $|M \cap \gamma| > o_M$ ,

and fix such an  $M$ . Now, let  $\eta$  be the least cardinal in  $M$  greater than  $o_M$ . Either  $\eta = o_M^+$  or  $\eta \cap M = o_M$ , so in particular  $|\eta \cap M| = o_M$ . By the minimality of  $\gamma$ ,  $|\gamma \cap M| = o_M^+$ . Since  $2^\gamma \in M$ , if  $o_M \in M$  then we have the following version of Chang's Conjecture:  $\langle \gamma, o_M^+ \rangle \rightarrow \langle o_M^+, o_M \rangle$ . To see this, note that otherwise there would be a function  $F: [\gamma]^{<\omega} \rightarrow \gamma$  for which there exists no set  $Z \subseteq \gamma$  closed under  $F$  with  $|Z| = o_M^+$  and  $|Z \cap o_M^+| = o_M$ . Then since  $\gamma$ ,  $o_M$  and  $o_M^+$  are in  $M$  there must be such a function in  $M$ , but since  $M \cap \gamma$  is closed under any function  $F': [\gamma]^{<\omega} \rightarrow \gamma$  in  $M$ , we have a contradiction. If  $o_M \notin M$ , we have the weaker statement that for every function  $F: [\gamma]^{<\omega} \rightarrow \gamma$  there exists a  $Z \subseteq \gamma$  closed under  $F$  such that  $\eta \geq |Z \cap \gamma| = |Z \cap \eta|^+$ . This weaker statement is implied by the version of Chang's Conjecture from the first case, and in fact is equivalent to the failure of (B). The failure of (B) then has consistency strength somewhere in between Chang's Conjecture and the existence of the sharp of every real (see [7]). In particular, (B) is weaker than the assumption “ $0^\#$  does not exist”, which has been used in other papers in the references, and (B) suffices for those arguments.

We will frequently be using the following consequence of (B):

**Lemma 1.6.** *Assume (B) and suppose  $\phi$  is a cardinal function on topological spaces such that  $\phi(X)$  is bounded by some finite iteration of the exponential function applied to  $|X|$ , and  $|\phi(X) \cap M| \geq \phi(X_M)$ . Then  $M \supseteq \phi(X_M)$ .*

*Proof.* Suppose  $\phi(X) \leq 2^{2^{\dots^{2^{|X|}}}}$ . By (B), it suffices to show  $|\phi(X) \cap M| \geq \phi(X_M)$  which we have assumed.  $\square$

## 2. EXAMPLES

The following example shows that we can have, modulo an inaccessible, two different elementary submodels  $M$  and  $N$  of the same size and a compact space  $X$  such that  $X_M$  is compact but  $X_N$  is not compact. We do not think the inaccessible is necessary, but we do not have another example.

**Example 2.1.** *A space  $X$  and elementary submodels  $M$  and  $N$  containing  $X$  such that  $|M| = |N|$ ,  $X_M$  is compact, but  $X_N$  is not.*

Let  $\kappa$  be an inaccessible cardinal smaller than the first 1-extendible. Take  $X$  to be the one-point compactification of the disjoint sum of  $D^\gamma$ , for  $\gamma < \kappa$ . Let  $M$  be an elementary submodel of a suitable  $H(\theta)$  with the property that  $M \cap \kappa$  is an ordinal less than  $\kappa$ , and such that all subsets of  $\gamma$  are in  $M$  whenever  $\gamma \in M \cap \kappa$ . Then  $X_M$  is compact. To get such an  $M$ , build an increasing sequence of elementary submodels  $M_n$  ( $n < \omega$ ) of cardinality less than  $\kappa$  such that for each even  $n$ ,  $M_n \cap \kappa$  is an ordinal, and such that for each odd  $n$ ,  $M_n$  contains the powerset of  $\gamma$  whenever  $\gamma \in M_k \cap \kappa$ , for  $k < n$ . Then  $\bigcup_{n \in \omega} M_n$  will be as desired. Note that for this construction to work we need  $\kappa$  to be a strong limit, but we also need  $\kappa$  to be regular: if not, we would have  $cf \kappa \in M$ , and therefore  $cf \kappa \subseteq M$ ; but then a cofinal subset of  $\kappa$  would be in  $M$ , so  $M \cap \kappa$  would have to be  $\kappa$ .

Now take  $N$  to be another elementary submodel such that  $|N| = \kappa$ ,  $X \in M$  and  $\kappa \in N$ . Then  $2^\kappa \in N$  but  $2^\kappa \not\subseteq N$ . Therefore,  $X_N$  is not compact, since (by Lemma 1.2)  $D^\kappa$  is not squashable and is not included in  $N$ , yet  $(D^\kappa)_N$  is a closed subspace of  $X_N$ .

We know that compact scattered spaces are squashable [6]. It is easy to get examples of non-scattered spaces that are squashable, like the previous one, taking perfect pre-images of scattered spaces in the correct way. (A map is *perfect* if it is continuous, closed, and points have compact inverses.) However not all squashable spaces are like that, even assuming there are no large cardinals.

**Example 2.2.** Let  $X$  be the long closed interval of length  $\kappa + 1$ ,  $\kappa > 2^{\aleph_0}$ . Then  $X$  is a connected squashable space – just pick  $M$  countably closed such that  $|M| < \kappa$ . Since  $X$  is connected,  $X$  cannot be a perfect pre-image of a scattered space.

**Problem.** Assuming say (B), characterize topologically the class of squashable spaces.

### 3. $\kappa$ -SCATTERED AND STRONGLY $\kappa$ -SCATTERED SPACES

We shall look at two generalizations of scattered spaces:

**Definition 3.1.** For  $p \in F \subseteq X$ ,  $\chi(p, F)$  is the least cardinality of a neighbourhood base for  $p$  in the subspace  $F$ .  $\chi(X) = \sup \{\chi(p, X) : p \in X\}$ .  $\chi(F, X)$  is the least cardinality of a neighbourhood base about  $F$  in  $X$ .  $\pi\chi(x, X)$  is the least cardinality of a collection of non-empty open sets such that every open set about  $x$  includes one.  $\pi\chi(X) = \sup \{\pi\chi(x, X) : x \in X\}$ . Clearly  $\pi\chi(X) \leq \chi(X)$ .

**Definition 3.2.** [2]. A space  $X$  is  $\kappa$ -scattered if for every closed subset  $F$  of  $X$ , there is a  $p \in F$  such that  $\chi(p, F) < \kappa$ .

**Definition 3.3.** A space  $X$  is *strongly*  $\kappa$ -scattered if for every closed subset  $F$  of  $X$ , there is a  $p \in F$  and an neighbourhood  $V$  of  $p$  such that  $|V \cap F| < \kappa$ .

In [6] it is shown that if  $X$  is compact and scattered, then  $X_M$  is compact. One could hope to generalize this result to  $\kappa$ -scattered, assuming maybe that  $\kappa \subseteq M$ . But this is consistently not true:

**Example 3.4.** Let  $\kappa$  be a cardinal and suppose  $\kappa^+ < 2^\kappa$ . Let  $\theta$  be a regular cardinal greater than  $2^\kappa$ , and let  $M$  be an elementary submodel of  $H(\theta)$  of cardinality  $\kappa^+$  including  $\kappa^+ + 1$ . Then  $(D^\kappa)_M$  is not compact. Let  $X$  be the one-point compactification of  $\kappa^+$  disjoint copies of  $D^\kappa$ . Then  $X$  is  $\kappa^+$ -scattered, but  $X_M$  is not compact.

However we can still improve some results in [6].

If  $X$  is any topological space and  $\leq_X$  is a well-ordering of a basis for  $X$ , then the *minimal decomposition* of  $X$  according to  $\leq_X$  is the sequence  $\langle P_\alpha, O_\alpha, X_\alpha : \alpha < \gamma \rangle$  defined as follows. For each  $\alpha < \gamma$ , let  $X_\alpha = X \setminus \bigcup \{O_\beta : \beta < \alpha\}$  (so  $X_0 = X$ ; the construction continues as long as the  $X_\alpha$ 's are non-empty), let  $O_\alpha$  be the  $\leq_X$ -least member of the basis for  $X$  such that  $O_\alpha \cap X_\alpha$  is non-empty and has the smallest possible cardinality, and let  $P_\alpha$  be  $O_\alpha \cap X_\alpha$ . If  $X$  is compact, this construction must end at a successor stage (and so we write  $\langle P_\alpha, O_\alpha, X_\alpha : \alpha \leq \gamma \rangle$ ).

**Theorem 3.5.** *Let  $X$  be a compact space, let  $\theta$  be a regular cardinal greater than  $2^{|X|}$  and let  $M$  be an elementary submodel of  $H(\theta)$ . Let  $\leq_X$  be a well-ordering in  $M$  of a basis for  $X$ , and let  $\langle P_\alpha, O_\alpha, X_\alpha : \alpha \leq \gamma \rangle$  be the minimal decomposition of  $X$  according to  $\leq_X$ . If  $M$  is closed under sequences of length  $|P_\alpha|$  for each  $\alpha \in (\gamma + 1) \cap M$ , then  $X_M$  is compact.*

*Proof.* Let  $D$  be an open cover of  $X_M$ , and let  $\gamma^*$  be the least  $\alpha \in (\gamma + 1) \cap M$  such that  $(X_{\gamma^*})_M$  can be covered by a finite subcover  $D_0$  of  $D$ . Then  $X \setminus \bigcup D_0$  is compact, and  $D_0$  is in  $M$ , so  $\gamma^*$  must be a successor ordinal or 0. Towards a contradiction, suppose that  $\gamma^* = \eta^* + 1$ . Now,  $X_{\eta^*} \setminus \bigcup D_0$  is compact, and since  $X_{\eta^*} \setminus \bigcup D_0 \subseteq P_{\eta^*}$ ,  $X_{\eta^*} \setminus \bigcup D_0 = (X_{\eta^*} \setminus \bigcup D_0)_M$ . Therefore, some finite subcover  $D_1$  of  $D$  covers  $(X_{\eta^*} \setminus \bigcup D_0)_M$ . Then  $D_0 \cup D_1$  is a finite subcover of  $D$  covering  $(X_{\eta^*})_M$ , giving a contradiction.  $\square$

Theorem 3.5 has the following corollary.

**Corollary 3.6.** *If  $M$  is  $\kappa$ -closed (i.e. subsets of  $M$  of size  $\leq \kappa$  are in  $M$ ) and  $X$  is strongly  $\kappa^+$ -scattered and compact, then  $X_M$  is compact.*

We next relate characters to squashability. A key concept in Kunen's work [8] is the following:

**Definition 3.7.** A  $\lambda$ -Čech-Pospíšil tree in a space  $X$  is a tree  $\mathcal{K} = \{K_s : s \in {}^{\leq \lambda}2\}$  satisfying:

- i) Each  $K_s$  is non-empty and closed in  $X$ ;
- ii)  $s \subseteq t$  implies  $K_s \supseteq K_t$ ;
- iii)  $K_{s0} \cap K_{s1} = \emptyset$ ;
- iv) if the length of  $s$  is  $\gamma$ , a limit ordinal, then  $K_s = \bigcap_{\alpha < \gamma} K_{s \upharpoonright \alpha}$ .

Čech and Pospíšil proved (see e. g. [3, 3.16]):

**Lemma 3.8.** *If  $X$  is compact and for each  $x \in X$ ,  $\chi(x, X) \geq \lambda$ , then there is a  $\lambda$ -Čech-Pospíšil tree in  $X$ , and hence  $|X| \geq 2^\lambda$ .*

We have the following results. We first use a proof from [8] to show:

**Lemma 3.9.** *If  $X_M$  is compact,  $\kappa + 1 \subseteq M$  and  $\chi(x, X) \geq \kappa$ , for every  $x \in X$ , then  $2^\kappa \subseteq M$ .*

*Proof.* Let  $\{K_s : s \in {}^{\leq \kappa}2\}$  be a  $\kappa$ -Čech-Pospíšil tree in  $X$ . By elementarity, we can suppose it is in  $M$ . We will prove by induction that  $2^\gamma \subseteq M$ , for every ordinal  $\gamma \leq \kappa$ . If  $\gamma$  is a successor ordinal, this is immediate. So suppose  $\gamma$  is a limit ordinal. Note that  $\gamma \in M$  since  $\kappa + 1 \subseteq M$ . Fix  $s \in 2^\gamma$ . By the induction hypothesis, we have  $s \upharpoonright \alpha \in M$ , for every  $\alpha < \gamma$ . Thus  $K_{s \upharpoonright \alpha} \in M$ . Also, by elementarity,  $(K_{s \upharpoonright \alpha})_M$  is closed in  $X_M$ , for every  $\alpha < \gamma$ . Since  $X_M$

is compact and  $\{(K_{s|\alpha})_M : \alpha < \gamma\}$  is centered, we have that there is an  $x \in \bigcap_{\alpha < \gamma} (K_{s|\alpha})_M$ . Note that  $x \in M$ , and therefore

$$s = \bigcup \{t : x \in K_t \text{ and } \text{length}(t) < \gamma\} \in M,$$

and we are done.  $\square$

By induction, we can now get as far as the first inaccessible, without assuming  $\kappa \subseteq M$ .

**Theorem 3.10.** *If  $X_M$  is compact,  $\kappa \in M$  is less than the first inaccessible, and  $\chi(x, X) \geq \kappa$  for every  $x \in X$ , then  $2^\kappa \subseteq M$ .*

*Proof.* Let  $\kappa$  be the least counterexample. By the previous lemma,  $\kappa$  cannot be  $\aleph_0$ . Also,  $\kappa$  cannot be  $\lambda^+$ , else  $\lambda \in M$  and hence  $2^\lambda \subseteq M$ , so  $\kappa \subseteq M$  and we can apply the previous lemma. A similar argument gives us that  $\kappa$  cannot be a limit cardinal that is not a strong limit. Finally, if  $\text{cf}(\kappa) = \lambda < \kappa$ , then  $\lambda \in M$  and there is a sequence  $\{\lambda_\alpha\}_{\alpha < \lambda}$  of cardinals in  $M$  with supremum  $\kappa$ . By the minimality of  $\kappa$ , we have that  $\lambda$  is included in  $M$  and that each  $2^{\lambda_\alpha} \subseteq M$ . Since  $\kappa$  is the supremum of the  $2^{\lambda_\alpha}$ 's, we have that  $\kappa \subseteq M$ , so by the previous lemma,  $2^\kappa \subseteq M$ .  $\square$

We suspect that 3.10 can be improved, replacing “inaccessible” by “1-extendible”, but we have been unable to prove that.

The following result from [6] will be useful:

**Lemma 3.11.** *If  $\chi(X) \leq \kappa \subseteq M$  and  $X_M$  is compact, then  $X = X_M$ .*

From this we deduce:

**Theorem 3.12.** *Assume (B). Suppose that a topological space  $X$  is squashed by an elementary submodel  $M$  such that  $|\mathcal{P}(X) \cap M| \in M$ . Then there is an  $x \in X$  such that  $\chi(x, X) < |\mathcal{P}(X) \cap M|$  and there is a  $y \in X$  such that  $\chi(y, X) > |\mathcal{P}(X) \cap M|$ .*

*Proof.* By (B),  $|\mathcal{P}(X) \cap M| = |X \cap M| = o_M$ . If  $\chi(x, X) \geq o_M$ , for every  $x \in X$ , then by the definition of  $o_M$ ,  $o_M \subseteq M$  and thus by Lemma 3.9, we would have  $2^{o_M} \subseteq M$ , a contradiction. If  $\chi(x, X) \leq o_M$ , for every  $x \in X$ , by Lemma 3.11, we would have that  $X_M$  is not compact, also a contradiction.  $\square$

Using 3.9, we can also show:

**Theorem 3.13.** *Assume (B). Suppose that a topological space  $X$  is squashed by an elementary submodel  $M$  such that  $|\mathcal{P}(X) \cap M| \in M$ . Then  $X$  is  $|\mathcal{P}(X) \cap M|$ -scattered.*

*Proof.* Fix  $X$  and  $M$  as in the statement of the theorem. Let  $\kappa$  denote  $|\mathcal{P}(X) \cap M|$ . By (B),  $o_M = \kappa \subseteq M$ . Suppose that  $X$  is not  $\kappa$ -scattered. Then there is a closed subset  $F$  of  $X$  such that  $\chi(x, F) \geq \kappa$ , for every  $x \in F$ . By elementarity we can take  $F \in M$  and we will also have that  $F_M$  is a closed subspace of  $X_M$ . Since  $X_M$  is compact,  $F_M$  will also be compact. Now using Lemma 3.9 for  $F$ , we would have  $2^\kappa \subseteq M$ , a contradiction.  $\square$

By Example 2.2, the hypothesis  $|\mathcal{P}(X) \cap M| \in M$  cannot be removed in the previous theorems. Since  $\omega$  is included in every elementary submodel of every  $H(\theta)$ , the same proof shows the following result from [6]:

**Corollary 3.14.** *If  $X$  is compact and squashed by a countable elementary submodel, then  $X$  is scattered.*

The following cardinal functions will be useful now and later:

**Definition 3.15.**  $c(X)$  is the sup of cardinalities of disjoint collections of open sets.  $d(X)$  is the least cardinality of a dense subset of  $X$ .  $\pi w(X)$  is the least cardinality of a collection  $\mathcal{P}$  of non-empty open sets such that each non-empty open set includes a member of  $\mathcal{P}$ .  $w(X)$  is the least cardinality of a basis for  $X$ . Clearly  $c(X) \leq d(X) \leq \pi w(X) \leq w(X)$ .

Now we quote the following result from [12]:

**Lemma 3.16.** *Suppose  $X_M$  is compact and either  $\chi(X_M) \leq \lambda$  or  $d(X_M) \leq \lambda$ . If  $2^\lambda \subseteq M$ , then  $X = X_M$ .*

We will also need the following result from [4]:

**Lemma 3.17.** *If  $X_M$  is compact, then there is a perfect map from  $X$  onto  $X_M$ , and hence  $X$  is compact.*

In investigating whether or not a compact space is squashable, a natural dichotomy occurs between the  $\kappa$ -scattered and non- $\kappa$ -scattered cases. We will first consider the non- $\kappa$ -scattered case.

**Theorem 3.18.** *Suppose  $\kappa \in M$  is less than the first inaccessible cardinal or suppose  $\kappa + 1 \subseteq M$ . Suppose  $X_M$  is compact and  $X$  (or  $X_M$ ) is not  $\kappa$ -scattered. If  $\chi(X_M) \leq \kappa$  or  $d(X_M) \leq \kappa$ , then  $X_M = X$ .*

*Proof.* We first deal with the case when  $X$  is not  $\kappa$ -scattered. Then there is a closed  $F \subseteq X$ ,  $F \in M$ , such that  $\chi(p, F) \geq \kappa$  for every  $p \in F$ . By 3.10 or by 3.9,  $2^\kappa \subseteq M$ , and by 3.16,  $X = X_M$ .

Now suppose instead that  $X_M$  is not  $\kappa$ -scattered. Then by 3.8, there is a  $\kappa$ -Čech-Pospíšil tree in  $X_M$ . Pulling back via the perfect map, we get a  $\kappa$ -Čech-Pospíšil tree in  $X$ , hence by the proof of 3.9 and 3.10, we again get  $2^\kappa \subseteq M$ .  $\square$

If we assume (B), we do not have to worry about inaccessibility provided our knowledge of  $X_M$ 's cardinal functions is sharp:

**Theorem 3.19.** *Assume (B). Suppose  $X_M$  is compact and  $\chi(X_M) = \kappa$  or  $d(X_M) = \kappa$  or  $\pi w(X_M) = \kappa$ , for some  $\kappa \in M$ . If either  $X$  or  $X_M$  is not  $\kappa$ -scattered, then  $X = X_M$ .*

*Proof.* Assuming (B), by the previous theorem, we just have to show that  $\kappa \subseteq M$ . To see this, it suffices to note that since  $X$  is compact,  $\chi(X) \leq |X|$ , so we can apply 1.6, since  $|\chi(X) \cap M| \geq \chi(X_M) = \kappa$ . A similar argument works for  $d$  or for  $\pi w$ .  $\square$

The following structure lemma for  $\kappa$ -scattered compact spaces will be used in the next two sections. It slightly strengthens a result of Efimov [2].

**Lemma 3.20.** *Suppose  $X$  is a  $\kappa$ -scattered compact space with  $\pi w(X) = \kappa$ ,  $cf(\kappa) = \omega$ ,  $\kappa > \omega$ . Then for any increasing sequence of cardinals  $\{\kappa_n\}_{n < \omega}$ , with  $\sup_{n < \omega} \kappa_n = \kappa$ ,  $\kappa_n$  regular, there exist regular closed subspaces  $\{X_n\}_{n < \omega}$  of  $X$  such that:*

- (a)  $\bigcup_{n < \omega} X_n$  is dense in  $X$ ;
- (b)  $\{y \in X_n : \chi(y, X) < \kappa_n\}$  is dense in  $X_n$ ;
- (c)  $\pi w(X) = \sum_{n \in \omega} \pi w(X_n)$ .

*Proof.* Fix an increasing sequence of cardinals  $\{\kappa_n : n \in \omega\}$  as in the hypothesis and define

$$E_n = \{x \in X : \chi(x, X) < \kappa_n\}.$$

First note that since  $X$  is  $\kappa$ -scattered, then  $D = \{x \in X : \chi(x, X) < \kappa\}$  is  $G_\delta$ -dense in  $X$  (see e.g. [12]). We give the proof here for completeness. Let  $V$  be a non-empty  $G_\delta$  set. Then there is a non-empty closed  $G_\delta$  set  $F \subseteq V$ . Since  $X$  is  $\kappa$ -scattered, there is an  $x \in F$  such that  $\chi(x, F) < \kappa$ . But then

$$\chi(x, X) \leq \chi(x, F) \cdot \chi(F, X) \leq \chi(x, F) \cdot \omega < \kappa.$$

Thus  $x \in D \cap V$  and we are done.

Let  $F_n = \overline{E_n}$ . Since  $D$  is  $G_\delta$ -dense in  $X$ , we have that  $X = \bigcup_{n \in \omega} F_n$ . Indeed, if  $X \setminus \bigcup_{n \in \omega} F_n = \bigcap_{n \in \omega} X \setminus F_n \neq \emptyset$ , then it would have to intersect  $D$ , a contradiction. Define  $X_n = \overline{\text{int} F_n}$ . Note that  $X_n \supseteq \overline{\text{int} X_n} \supseteq \overline{\text{int} F_n} = X_n$ , so  $X_n$  is a regular closed subspace of  $X$ . By the Baire Category Theorem, some  $F_n$ , hence all  $F_n$  from some  $n_0$  onward have non-empty interiors. It follows that  $\bigcup_{n \in \omega} \text{int} F_n$  and, a fortiori,  $\bigcup_{n \in \omega} X_n$  is dense in  $X$ . To see this, suppose there were a non-empty open  $V \subseteq X - \bigcup_{n < \omega} \text{int} F_n$ . Then  $V = \bigcup_{n < \omega} F_n \cap V$ . Again by Baire Category, for some  $n$ ,  $F_n \cap V$  has non-empty interior in  $V$  and hence in  $X$ . But  $\text{int}(F_n \cap V) \subseteq \text{int} F_n \subseteq X - V$ , contradiction. Also, since  $\pi$ -weight is inherited by and from dense sets,  $\pi w(X) = \pi w(\bigcup_{n \in \omega} X_n)$ , so  $\pi w(X) \leq \sum_{n \in \omega} \pi w(X_n)$ . On the other hand, since  $\pi w(\text{int} F_n) \leq \pi w(X)$ , we have  $\pi w(X) = \sum_{n \in \omega} \pi w(X_n)$ .

Note that since  $E_n$  is dense in  $F_n$ ,  $E_n \cap \text{int} F_n$  is dense in  $\text{int} F_n$  and hence in  $\overline{\text{int} F_n} = X_n$ . Thus  $X_n$  has a dense subspace of points of character  $< \kappa_n$ .  $\square$

#### 4. THE COUNTABLE CHAIN CONDITION CASE

In [4], [6], [8], [10], [11], [12] it is shown that if  $X_M$  is compact and satisfies various properties stronger than the countable chain condition, then  $X = X_M$ , if  $|X|$  is small or anti-large-cardinal assumptions are made. It is therefore a natural question whether assuming  $X_M$  is compact and satisfies the countable chain condition is enough to get that  $X$  is not squashable. We will show this question is undecidable. First we have a consistent example.

**Example 4.1.** “ $\Psi$ -space” is any space obtained by taking a maximal almost disjoint family  $\mathcal{A}$  of subsets of  $\omega$ , and putting a topology on  $\mathcal{A} \cup \omega$  as follows. Each point in  $\omega$  is isolated. For each  $A \in \mathcal{A}$ , a neighbourhood of  $A$  is  $\{A\}$  union a cofinite piece of  $A$ .

Assume  $\neg CH$ . Then the one-point compactification  $X$  of  $\Psi$ -space satisfies the countable chain condition and is scattered and compact. Then  $X_M$  is compact, but it is different from  $X$  if  $|M| < |X|$ . So  $X$  is squashable.

**Problem.** Is there a consistent example without assuming large cardinals which is not scattered?

On the way to proving a consistent theorem, we first show a partial positive result:



**Theorem 4.2.** *Assume (B) and  $\kappa \in M$ . Suppose  $X_M$  is compact and satisfies the countable chain condition, and  $d(X_M) = \kappa$  or  $\chi(X_M) = \kappa$ , where  $\kappa^\omega = \kappa$ . Then  $X_M = X$ .*

The following lemma of Šapirovsĭiĭ (see e.g. [3]) will be useful.

**Lemma 4.3.**  $w(X) \leq \pi\chi(X)^{c(X)}$ .

We also need the following lemma proved in [6, Theorem 4.9] (there we have the assumption  $2^\omega \subseteq M$ , but just  $\omega_1 \subseteq M$  is used for this fact):

**Lemma 4.4.** *If  $c(X_M) = \aleph_0$  and  $\omega_1 \subseteq M$ , then  $c(X) = \aleph_0$ .*

*Proof of 4.2.* By 1.6 we have  $\kappa \subseteq M$ . Since  $\kappa^\omega = \kappa$ ,  $\kappa$  is uncountable, so by 4.4,  $X$  satisfies the countable chain condition. If  $X$  is not  $\kappa$ -scattered we are done by Theorem 3.19, so we can suppose  $X$  is  $\kappa$ -scattered. But then as we saw before,

$$D = \{x \in X : \pi\chi(x, X) < \kappa\}$$

is dense in  $X$ .

We will have that  $w(D) \leq \kappa^\omega = \kappa$ , since  $\pi\chi(D) \leq \kappa$  and  $D$  by density also satisfies the countable chain condition. But then  $\pi\chi(X) \leq \pi w(X) = \pi w(D) \leq w(D) \leq \kappa$  (since  $X$  is regular), whence by 4.3 again,  $w(X) \leq \kappa^\omega = \kappa$ .

Since  $\kappa \subseteq M$ , this implies that  $X_M = X$ . □

We can now show a consistent positive result. Let *SCH* stand for the *Singular Cardinals Hypothesis*, namely that for every singular cardinal  $\kappa$ , if  $2^{cf(\kappa)} < \kappa$ , then  $\kappa^{cf(\kappa)} = \kappa^+$ .

**Theorem 4.5.** *Assume (B) and  $CH + SCH$ . Then if  $X_M$  is uncountable and compact,  $\chi(X_M) \in M$  and  $X_M$  satisfies the countable chain condition, then  $X_M = X$ .*

*Proof.* Let  $\kappa = \chi(X_M)$ . The case when  $\kappa \leq \aleph_0$  was done in [6] in *ZFC*, so we may assume  $\kappa \geq \aleph_1$ . Then, as before, since we are assuming (B), we have  $\kappa \subseteq M$ . Therefore, by lemma 4.4,  $X$  satisfies the countable chain condition.

If  $X$  is not  $\kappa$ -scattered, then, again, we are done by Theorem 3.19. If  $cf(\kappa) > \omega$ , then by *CH + SCH* we have  $\kappa^\omega = \kappa$  (*CH* takes care of  $\kappa = \omega_1 = 2^\omega$  and *SCH* of the others), and therefore, by the previous theorem, we are also done.

Suppose then that  $X$  is  $\kappa$ -scattered and that  $cf(\kappa) = \omega$ . The proof uses ideas from [12].

Fix  $\{\kappa_n\}_{n \in \omega}$ ,  $E_n$  and  $X_n$  as in the proof of Lemma 3.20. It will then suffice to show each  $(X_n)_M = X_n$ . Note that each  $X_n$  satisfies the countable chain condition since they are regular closed subspaces of  $X$ . We can also assume that  $\kappa_0 > \omega_1$ .

Fix  $n \in \omega$ . To show that  $(X_n)_M = X_n$  we can now repeat the same argument done in the proof of 4.2. We know that  $E_n = \{x \in X : \chi(x, X) < \kappa_n\}$  is dense in  $X_n$  and that  $\pi\chi(E_n) \leq \chi(E_n) \leq \kappa_n$ . Since  $X_n$  satisfies the countable chain condition, so does  $E_n$  and therefore  $w(E_n) \leq \kappa_n^\omega = \kappa_n$ , by *CH + SCH* (since  $\kappa_n$  is regular and  $> \omega_1 = 2^\omega$ ). But then as before we can conclude that  $\pi\chi(X_n) \leq \kappa_n$  and therefore  $w(X_n) \leq \kappa_n^\omega = \kappa_n$ . Since  $\kappa \subseteq M$ , we have  $\kappa_n \subseteq M$  and we are done by 3.11. □

**Problem.** Is *CH + SCH* necessary? Are there *ZFC + CH + SCH* results below some large cardinal?

Note the hypothesis of  $X_M$  being uncountable is essential; otherwise just pick  $X$  to be any uncountable compact  $T_2$  scattered space and  $M$  any countable elementary submodel. By [6],  $X_M$  will be compact. Also it satisfies the countable chain condition, but  $X_M \neq X$ .

## 5. $X_M$ 'S CO-ABSOLUTE WITH DYADIC SPACES

**Definition 5.1.** Two spaces are *co-absolute* if they have isomorphic regular-open algebras. A compact space is *dyadic* if it is the continuous image of some power of the two-point discrete space.

In [2] it is shown that:

**Theorem 5.2.** *If  $\lambda$  is an infinite cardinal with  $cf(\lambda) \geq \omega_1$  and  $X$  is a compact space co-absolute with a dyadic one with  $\pi w(X) = \lambda$ , then  $X$  is not  $\lambda$ -scattered.*

We then have:

**Theorem 5.3.** *Assume (B) or that  $\kappa <$  the first inaccessible cardinal and  $\kappa \in M$ . If  $X_M$  is co-absolute with a dyadic compactum with  $\pi w(X_M) = \kappa$ , where  $cf(\kappa) \geq \omega_1$ , then  $X = X_M$ .*

*Proof.* If  $X_M$  is co-absolute with a dyadic compactum, then  $X_M$  is compact. By the previous theorem,  $X_M$  is not  $\kappa$ -scattered. Since  $d(X_M) \leq \pi w(X_M) = \kappa$ , by Theorem 3.18 or 3.19 we then have that  $X = X_M$ .  $\square$

To show the general result we will need the following lemmas:

**Lemma 5.4.** [9]. *If  $X$  is co-absolute with a dyadic compactum, then  $X$  is co-absolute with either a finite disjoint sum of powers of  $D$  or else with the one-point compactification of a countable disjoint sum of powers of  $D$ .*

**Lemma 5.5.** *Each regular closed subspace of a compact space co-absolute with a dyadic compactum is itself co-absolute with a dyadic compactum.*

*Proof.* Let  $X$  be a compact space co-absolute with  $K$ , a dyadic compactum. Let  $Z$  be a regular closed subspace of  $X$ . Let  $i$  be an isomorphism between the algebra of regular closed subspaces of  $X$  and the algebra of regular closed subspaces of  $K$ . Such an isomorphism exists, since the dual regular open algebras are isomorphic. Let  $i(Z) = L$ . By [1],  $L$  is dyadic, and clearly the algebra of regular closed subspaces of  $Z$  is isomorphic to the algebra of regular closed subspaces of  $L$ . It follows that  $Z$  and  $L$  are co-absolute.  $\square$

**Lemma 5.6.** [2]. *Suppose  $X$  is co-absolute to a dyadic compactum and  $X$  has a dense subspace of points of character, i.e.  $\chi(p, X)$ , less than  $\lambda$ , where  $\lambda$  is an uncountable regular cardinal. Then  $\pi w(X) < \lambda$ .*

Efimov [2] states this for “ $\delta$ -character” rather than character, but the former does not exceed the latter.

We are now ready to show our main result:

**Theorem 5.7.** *Assume (B) or that  $\kappa$  is less than the first inaccessible cardinal. Suppose  $X_M$  is compact and  $\pi w(X_M) = \kappa \in M$ , and  $X_M$  is co-absolute with a dyadic compactum. Then  $X_M = X$ .*

*Proof.* It follows from Lemma 5.4 that compact spaces co-absolute with dyadic compacta have no isolated points, and thus that if  $X_M$  is such a space, then  $2^{\aleph_0} \subseteq M$  [6]. Such  $X_M$ 's satisfy the countable chain condition; since  $\omega_1 \subseteq M$ , it follows that  $X$  does as well.

If  $X_M$  is not  $\pi w(X_M)$ -scattered, we are done by Theorem 3.19, so assume that it is. Then, by Theorem 5.2,  $\kappa = \pi w(X_M)$  has countable cofinality. Similarly assume  $X$  is  $\pi w(X_M)$ -scattered. We also may assume that  $\pi w(X_M)$  is uncountable, else  $X = X_M$  by Lemma 3.16, since  $2^{\aleph_0} \subseteq M$ .

Applying Lemma 3.20 to  $X$  and letting  $\{\kappa_n\}_{n < \omega}$  be a strictly increasing sequence of regular cardinals approaching  $\kappa$ , we can obtain a sequence of  $X_n$ 's. Without loss of generality, assume the sequence as well as the sequence of  $\kappa_n$ 's is in  $M$ . Then, by elementarity, we can get a sequence of  $X_n$ 's satisfying the following conditions:

- (d) each  $(X_n)_M$  is a regular closed subspace of  $X_M$ ;
- (e)  $\bigcup_{n < \omega} (X_n)_M$  is dense in  $X_M$ ;
- (f)  $\pi w(X_M) = \sum_{n < \omega} \pi w((X_n)_M)$ ;
- (g)  $\pi w((X_n)_M) < \kappa$ , for each  $n \in \omega$ .

To get (g), by Lemma 3.20 and elementarity, we have that for each  $n \in \omega$ ,  $\{y \in (X_n)_M : \chi(y, X_M) < \kappa_n\}$  is dense in  $(X_n)_M$ . Furthermore, note that for  $y \in (X_n)_M$ ,  $\chi(y, (X_n)_M) \leq \chi(y, X_n)$ . Applying Lemmas 5.5 and 5.6, we see that  $\pi w((X_n)_M) < \kappa_n < \kappa$ .

We claim that we can also obtain the  $X_n$ 's such that each  $(X_n)_M$  is not  $\pi w((X_n)_M)$ -scattered. Since  $\pi w((X_n)_M) < \kappa$ , by 5.2 and 5.5 the claim holds for  $\pi w(X_M) = \aleph_\omega$ . If the claim fails, there is a counterexample  $X$  with  $\pi w(X_M)$  minimal. Then none of the  $X_n$ 's, obtained as above, are counterexamples. Thus for each  $n$  we can obtain a sequence  $(Y_{nk})_{k \in \omega}$  satisfying all the conditions we want: for  $n$  such that  $X_n$  is already not  $\pi w((X_n)_M)$ -scattered, we just take  $Y_{nk} = X_n$  for each  $k$ ; for  $n$  such that  $X_n$  is  $\pi w((X_n)_M)$ -scattered, we apply Lemma 3.20 and our induction hypothesis. But then the set of all  $Y_{nk}$ 's for all  $X_n$ 's is the desired countable collection of subspaces of  $X$ .

Now, since  $(X_n)_M$  is not  $\pi w((X_n)_M)$ -scattered, for any  $n \in \omega$ , by the proofs of Theorem 3.18 and Theorem 3.19, we conclude that  $2^{< \kappa} \subseteq M$ , and so  $\kappa \subseteq M$ . Consider two sub-cases:  $\kappa$  is or is not a strong limit. In the former sub-case, the proof of Theorem 4.5 works without (B) and  $CH + SCH$  to get that  $X = X_M$ . In the latter sub-case, there is a regular  $\mu < \kappa$ , such that  $2^\mu = 2^\kappa$ . Thus  $2^\mu \subseteq M$ . Then by Lemma 3.16,  $X_M = X$ .  $\square$

## 6. BOOLEAN ALGEBRAS

One might expect that our topological results should have some implications for Boolean algebras, and they do. In [6] the following result was proved, where for a Boolean algebra  $A$ , " $S(A)$ " denotes the Stone space of  $A$ :

**Theorem 6.1.** *Assume  $0^\#$  does not exist. Let  $A$  be a Boolean algebra. If  $A \cap M$  is complete and  $(S(A))_M$  is compact, then  $A = A \cap M$ .*

Using our new topological results, we obtain:

**Theorem 6.2.** *Suppose (B) and that  $A$  is a Boolean algebra such that  $(S(A))_M$  is compact. If the completion of  $A \cap M$  is isomorphic to the completion of some Boolean algebra  $C$  homomorphically embedded in the clopen algebra  $K$  of some  $D^\kappa$ , then  $A = A \cap M$ . The conclusion holds in ZFC for  $A$ 's with  $|A|$  less than the first inaccessible.*

*Proof.* In [6] it was shown that if  $(S(A))_M$  is compact, then  $(S(A))_M = S(A \cap M)$ . To say that  $C$  is homomorphically embedded in  $K$  says that  $S(C)$  is a continuous image of  $D^\kappa$ . We are then assuming that  $(S(A))_M$  is co-absolute with  $S(C)$ , and that the latter is dyadic. Therefore  $(S(A))_M = S(A)$ . But then  $A = A \cap M$ .  $\square$

**Corollary 6.3.** *Suppose (B) and that  $A$  is a Boolean algebra such that  $S(A)_M$  is compact. If the completion of  $A \cap M$  is isomorphic to the algebra for adding  $\kappa$  many Cohen reals, then  $A = A \cap M$ . The conclusion holds in ZFC for  $A$ 's with  $|A|$  less than the first inaccessible.*

Assuming (B) and  $CH + SCH$ , we get a stronger result:

**Theorem 6.4.** *Suppose (B) and  $CH + SCH$  and that  $C$  is a Boolean algebra such that  $S(C)_M$  is compact and satisfies the countable chain condition. Then  $C = C \cap M$ .*

## REFERENCES

- [1] B. A. Efimov, *Dyadic bicompecta*, Trans. Moscow Math. Soc. **14** (1965), 229–267.
- [2] ———, *On the cardinality of compactifications of dyadic spaces*, Math. USSR Sbornik **25** (1975), no. 4, 579–593.
- [3] I. Juhász, *Cardinal functions in topology: ten years later*, Math. Centre Tracts, 1980.
- [4] L. R. Junqueira, *Upwards preservation by elementary submodels*, Top. Proc. **25** (Spring 2000), 225–249.
- [5] L. R. Junqueira and F. D. Tall, *The topology of elementary submodels*, Top. Appl. **82** (1998), 239–266.
- [6] ———, *More reflections on compactness*, Fund. Math. **176** (2003), 127–141.
- [7] A. Kanamori, *The higher infinite: large cardinals in set theory from their beginnings*, Springer-Verlag, Berlin, 1994.
- [8] K. Kunen, *Compact spaces, compact cardinals, and elementary submodels*, Top. Appl. **130** (2003), 99–109.
- [9] L. B. Shapiro, *On spaces coabsolute to dyadic compacta*, Sov. Math. Dokl. **35** (1987), 434–438.
- [10] F. D. Tall, *If it looks and smells like the reals....*, Fund. Math. **163** (2000), 1–11.
- [11] ———, *Reflecting on compact spaces*, Top. Proc. **25** (Summer 2000), 345–350.
- [12] ———, *Reflections on dyadic compacta*, Top. Appl. **137** (2004), 251–258.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSITY OF SÃO PAULO, CAIXA POSTAL 66281, SÃO PAULO, SP BRAZIL, 05315-970

*E-mail address:* lucia@ime.usp.br

DEPARTMENT OF MATHEMATICS, MIAMI UNIVERSITY, OXFORD, OHIO, UNITED STATES, 45056

*E-mail address:* larsonpb@muohio.edu

DEPT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA, M5S 3G3

*E-mail address:* tall@math.toronto.edu