

A CHOICE FUNCTION ON COUNTABLE SETS, FROM DETERMINACY

PAUL B. LARSON

ABSTRACT. We prove that $\text{AD}_{\mathbb{R}}$ implies the existence of a definable class function which, given a countable set X , a tall ideal I on ω containing Fin and a function from $I \setminus \text{Fin}$ to X which is invariant under finite changes, selects a nonempty finite subset of X . Among other applications, this gives an alternate proof of the fact (previously established by Di Prisco-Todorcevic [2]) that there is no selector for the E_0 degrees in the $\mathcal{P}(\omega)/\text{Fin}$ -extension of a model of $\text{AD}_{\mathbb{R}}$.

An ideal I on ω is said to be *tall* if every infinite subset of ω contains an infinite member of I . In this paper we show that, assuming the Axiom of Real Determinacy ($\text{AD}_{\mathbb{R}}$), there is a definable procedure which, given a tall ideal on ω containing Fin (the ideal of finite subsets of ω), and a function on $I \setminus \text{Fin}$ which is invariant under finite changes and has countable range, selects a nonempty finite subset of the range of this function. In the case we are most interested in, I is generated by a countable collection of pairwise orthogonal ideals. In this context, these ideals represent a $\mathcal{P}(\omega)/\text{Fin}$ -name for an element of a countable set from the ground model. Our result (Theorem 3.3) has the following consequence (Corollary 3.4) : if

- $M[U]$ is a $\mathcal{P}(\omega)/\text{Fin}$ -extension of a model M of $\text{AD}_{\mathbb{R}}$,
- F is a function in $M[U]$ with domain $Y \in M$,
- G is a function in M with domain Y such that, for all $y \in Y$, $G(y)$ is countable and $F(y)$ is a finite nonempty subset of $G(y)$,

then there exists a function G' in M with domain Y such that, for all $y \in Y$, $G'(y)$ is a nonempty finite subset of $G(y)$. This gives another proof of a theorem of Di Prisco-Todorcevic [2] (see also [6]) that there is no function in $M[U]$ which selects a single member from each E_0 -equivalence class (where E_0 is the relation of mod-finite agreement on $\mathcal{P}(\omega)$). Since the finite sets of E_0 degrees can be injectively embedded into the (individual) E_0 degrees in a Borel fashion, this also implies that whenever F is an equivalence relation on $\mathcal{P}(\omega)$ in M whose equivalence classes are countable unions of E_0 -classes, if a function in $M[U]$ picks an E_0 subclass for each F -class, then such a function exists already in M .

In the final section of the paper (Section 5) we note that “finite” cannot be replaced with “size n ” in our result, for any positive $n \in \omega$. In particular, since $\text{AD}_{\mathbb{R}}$ implies that all sets of reals have the property of Baire, for any fixed positive

Research supported by NSF Grants DMS-0801009 and DMS-1201494. Results obtained in February 2011. The author thanks Andrés Caicedo and Grigor Sargsyan for help with bibliographic references. This version of the paper corrects minor errors in the published version.

$n \in \omega$ it is not possible in this context to pick n members from each countable family of orthogonal ideals generating a tall ideal.

1. STRATEGICALLY SELECTIVE IDEALS

Given a sequence $\langle x_n : n \in \omega \rangle$ consisting of subsets of ω , a *fast selector* for this sequence is a set $\{b_i : i \in \omega\}$ (listed in increasing order) such that $b_0 \in x_0$ and $b_{i+1} \in x_{b_i}$ for all $i \in \omega$. Given an ideal I on ω , we let $G(I)$ be the game where players *Noselector* (who moves first) and *Selector* alternately pick the members of a \subseteq -decreasing sequence of I -positive sets, with *Selector* winning if the sequence produced has a fast selector in I^+ . We say that an ideal I is *strategically selective* if *Noselector* does not have a winning strategy in $G(I)$. The proof of Theorem 1.1 below consists of combining work of Martin, Solovay and Woodin with Mathias's proof [4] that $\text{AD}_{\mathbb{R}}$ implies that there are no tall selective ideals.

Theorem 1.1 ([3]). *If $\text{AD}_{\mathbb{R}}$ holds, then there are no tall strategically selective ideals on ω .*

It follows that under $\text{AD}_{\mathbb{R}}$, the strategically selective ideals are exactly those ideals I such that every infinite $A \subseteq \omega$ contains an infinite $B \subseteq \omega$ with $\mathcal{P}(B) \cap I$ containing no infinite set.

For the definition of strategically selective ideals, it is appropriate for *Noselector* to move first, but for the purposes of this paper *Selector* moves first in the relevant game. We therefore let $G'(I)$ be modified game in which *Selector* moves first, and again the two players alternately pick the members of a \subseteq -decreasing sequence of I -positive sets, with *Selector* winning if the sequence produced has a fast selector in I^+ . Theorem 1.1 implies that under $\text{AD}_{\mathbb{R}}$, *Noselector* still has a winning strategy in this game whenever I is a tall ideal on ω .

Corollary 1.2. *If $\text{AD}_{\mathbb{R}}$ holds, then *Noselector* has a winning strategy for $G'(I)$ whenever I is a tall ideal on ω .*

2. EMBEDDING THE BAIRE SPACE

For each \subseteq -decreasing sequence $\bar{x} = \langle x_n : n \in \omega \rangle$ consisting of infinite subsets of ω , we recursively define an association of finite sequences $\sigma \in \omega^{<\omega}$ to finite subsets $z_{\sigma}^{\bar{x}} \subseteq \omega$ as follows.

- $z_{\langle \rangle}^{\bar{x}} = \emptyset$;
- $z_{\sigma \smallfrown \langle 0 \rangle}^{\bar{x}} = z_{\sigma}^{\bar{x}}$;
- if σ has length n , i is an element of ω , and $z_{\sigma}^{\bar{x}} = \emptyset$, then $z_{\sigma \smallfrown \langle i+1 \rangle}^{\bar{x}}$ is the singleton containing the i th member of x_n , where the least member is taken to be the 0th member.
- if σ has length n and i is an element of ω , $z_{\sigma}^{\bar{x}}$ is nonempty, and r is the largest member of $z_{\sigma}^{\bar{x}}$, then $z_{\sigma \smallfrown \langle i+1 \rangle}^{\bar{x}}$ is the union of $z_{\sigma}^{\bar{x}}$ with the singleton containing the i th member of x_{r+n} greater than all elements of $z_{\sigma}^{\bar{x}}$.

Let $\pi_{\bar{x}}$ be the continuous embedding of the Baire space into $\mathcal{P}(\omega)$ induced by the assignment $\sigma \mapsto z_{\sigma}^{\bar{x}}$. Comeagerly many members of the Baire space map to infinite sets, and all of these are fast selectors for $\langle x_n : n \in \omega \rangle$.

2.1 Remark. Given \bar{x} as above, and $\sigma_0, \sigma_1 \in \omega^{<\omega}$, there exist $\tau_0, \tau_1 \in \omega^{<\omega}$ such that $\sigma_0 \widehat{\ } \tau_0$ and $\sigma_1 \widehat{\ } \tau_1$ have the same length, and $z_{\sigma_0 \widehat{\ } \tau_0}^{\bar{x}}$ and $z_{\sigma_1 \widehat{\ } \tau_1}^{\bar{x}}$ have the same maximal element. Furthermore, given $\rho_0, \rho_1 \in \omega^{<\omega}$ with the same length, if k is the maximal element of both $z_{\rho_0}^{\bar{x}}$ and $z_{\rho_1}^{\bar{x}}$, then for all $w \in \omega^\omega$,

$$\pi_{\bar{x}}(\rho_0 \widehat{\ } w) \setminus k = \pi_{\bar{x}}(\rho_1 \widehat{\ } w) \setminus k.$$

These observations are used in the first paragraph of the proof of Theorem 3.2.

2.2 Definition. Let $\bar{x} = \langle x_n : n \in \omega \rangle$ and $\bar{y} = \langle y_n : n \in \omega \rangle$ be \subseteq -decreasing sequences of infinite subsets of ω . We say that \bar{x} and \bar{y} are *eventually intertwining* if there exist $n, m \in \omega$ such that for all $k \in \omega$,

$$y_{m+k+1} \subseteq x_{n+k+1} \subseteq y_{m+k} \subseteq x_{n+k}.$$

The family of embeddings $\pi_{\bar{x}}$ was defined in order to make the following lemma hold.

Lemma 2.3. *Let \bar{x} and \bar{y} be eventually intertwining sequences of infinite subsets of ω , and let C be a comeager subset of the Baire space. Then there exist $a, b \in C$ such that $\pi_{\bar{x}}(a) = \pi_{\bar{y}}(b)$.*

Proof. We recursively define σ_n, τ_n in $\omega^{<\omega}$ for $n \in \omega$ as follows, maintaining the following conditions for all $n \in \omega$:

- $\sigma_n \subseteq \sigma_{n+1}$;
- $\tau_n \subseteq \tau_{n+1}$;
- $z_{\sigma_n}^{\bar{x}} = z_{\tau_n}^{\bar{y}}$.

Let $n_0, m_0 \in \omega$ witness that \bar{x} and \bar{y} are eventually intertwining. Let σ_0 consist of n_0 many 0's followed by $a+1$, and let τ_0 consist of m_0 many 0's, followed by $b+1$, where the a -th member of x_{n_0} is the same as the b -th member of y_{m_0} , and greater than both n_0 and m_0 . Suppose now that $n \in \omega$, σ_n and τ_n have been chosen. We need to see that, given a dense set $D \subseteq \omega^{<\omega}$, we can choose σ_{n+1} and τ_{n+1} as above so that $\sigma_{n+1} \in D$ (we need to do this also for the τ_n 's, but the argument is symmetric). If necessary, we first extend σ_n with 0's to a sequence σ'_n of length at least $|\tau_n| + (n_0 - m_0 + 1)$. Let σ_{n+1} be any extension of σ'_n in D . Suppose that $\langle i_0, \dots, i_p \rangle$ is the sequence added to σ'_n to make σ_{n+1} . We can add a sequence $\langle j_0, \dots, j_p \rangle$ to τ_n to make τ_{n+1} , in such a way that for all $q \leq p$,

$$z_{(\sigma'_n) \widehat{\ } \langle i_0, \dots, i_q \rangle}^{\bar{x}} = z_{\tau_n \widehat{\ } \langle j_0, \dots, j_q \rangle}^{\bar{y}}.$$

For each $q \leq p$, let $j_q = 0$ if and only if $i_q = 0$. Given $q \leq p$ such that $i_q > 0$, $z_{(\sigma'_n) \widehat{\ } \langle i_0, \dots, i_q \rangle}^{\bar{x}}$ has length at least 2. Let $r < s$ be its top two elements. Then s is in $x_{|\sigma'_n|+q+r}$, and therefore in $y_{|\tau_n|+q+r}$. It follows then that j_q can be chosen as desired, so that s is the $(j_q - 1)$ -th member of $y_{|\tau_n|+q+r}$ greater than all elements of $z_{(\sigma'_n) \widehat{\ } \langle i_0, \dots, i_{q-1} \rangle}^{\bar{x}}$. \square

3. EXTENDING A FUNCTION ON AN IDEAL

3.1 Definition. Let I be an ideal on ω containing Fin, and let \mathcal{J} be a set of ideals on ω . We say that the members of \mathcal{J} are *pairwise I -orthogonal* if $x \cap y \in I$ whenever J_0 and J_1 are distinct elements of \mathcal{J} , and x and y are elements of J_0 and J_1 respectively. We let $\text{FCI}(I, \mathcal{J})$ (the *finite covering ideal*) denote the ideal

generated by $I \cup \bigcup \mathcal{J}$. For each $x \in \text{FCI}(I, \mathcal{J}) \setminus I$, the set of $J \in \mathcal{J}$ such that $x \cap y \in I^+$ for some $y \in J$ is finite and nonempty. We call this set $C(I, \mathcal{J}, x)$ below.

The following is our main result. In Theorems 3.2 and 3.3, the formula defining the specified operation can be extracted from the proof, but we leave this to the interested reader.

Theorem 3.2. *Suppose that $\text{AD}_{\mathbb{R}}$ holds. There exists a definable operation which takes in a nonempty countable set X , a tall ideal I on ω containing Fin and a function*

$$f: (I \setminus \text{Fin}) \rightarrow ([X]^{<\omega} \setminus \{\emptyset\})$$

which is invariant under finite changes, and returns a nonempty set $X' \subseteq [X]^{<\omega}$ and a collection of pairwise I -orthogonal ideals $\{J_a : a \in X'\}$ such that each J_a properly contains I and every I -positive set contains an I -positive set in some J_a .

Proof. By Corollary 1.2, *Noselector* has a winning strategy Γ in the game $G'(I)$. Suppose now that \bar{x} is the sequence of moves played by *Noselector* in some run of $G'(I)$ which has been won by *Noselector*. The function $f \circ \pi_{\bar{x}}$ maps a comeager subset of the Baire space into a countable set. Since all sets of reals have the Baire property, and f is invariant under finite changes, $f \circ \pi_{\bar{x}}$ is constant on a comeager set (not merely a relatively comeager set; see Remark 2.1). Let $v_{\bar{x}}$ be this constant value.

Noselector can win the modified game in which he must declare with his first move the value $v_{\bar{x}}$ corresponding to the eventual run of the game. To see this, note that otherwise there exist a first move w for *Selector* and a sequence of $G'(I)$ strategies Σ_a for *Selector*, for each nonempty $a \in [X]^{<\omega}$, each guaranteeing that either *Selector* wins the game or that the value $v_{\bar{x}}$ will not be a . Fixing some enumeration $\langle a_i : i \in \omega \rangle$ of $[X]^{<\omega} \setminus \{\emptyset\}$, consider then the sequence which blends each Σ_{a_i} against Γ , i.e.,

- $x_0 = \Gamma(\langle w \rangle)$
- $y_0^0 = \Sigma_{a_0}(\langle x_0 \rangle)$
- $x_1 = \Gamma(\langle w, x_0, y_0^0 \rangle)$
- $y_1^0 = \Sigma_{a_0}(\langle x_0, y_0^0, x_1 \rangle)$
- $y_0^1 = \Sigma_{a_1}(\langle y_1^0 \rangle)$
- $x_2 = \Gamma(\langle w, x_0, y_0^0, x_1, y_0^1 \rangle)$
- $y_2^0 = \Sigma_{a_0}(\langle x_0, y_0^0, x_1, y_1^0, x_2 \rangle)$
- $y_1^1 = \Sigma_{a_1}(\langle y_1^0, y_0^1, y_2^0 \rangle)$
- $y_0^2 = \Sigma_{a_2}(\langle y_1^1 \rangle)$
- $x_3 = \Gamma(\langle w, x_0, y_0^0, x_1, y_0^1, x_2, y_0^2 \rangle)$
- $y_3^0 = \Sigma_{a_0}(\langle x_0, y_0^0, x_1, y_1^0, x_2, y_2^0, x_3 \rangle)$
- $y_2^1 = \Sigma_{a_1}(\langle y_1^0, y_0^1, y_2^0, y_1^1, y_3^0 \rangle)$
- $y_1^2 = \Sigma_{a_2}(\langle y_1^1, y_0^2, y_2^1 \rangle)$
- $y_0^3 = \Sigma_{a_3}(\langle y_1^2 \rangle)$
- $x_4 = \Gamma(\langle w, x_0, y_0^0, x_1, y_0^1, x_2, y_0^2, x_3, y_0^3 \rangle)$

and so on. Then the sequence $\bar{x} = \langle x_n : n \in \omega \rangle$ is the sequence of moves for *Noselector* in a run of $G'(I)$ according to Γ , and thus winning for *Noselector*. Letting i be such that $a_i = v_{\bar{x}}$, let \bar{y} be \bar{x} in the case where $i = 0$ and

$$\langle y_{n+1}^{i-1} : n \in \omega \rangle$$

otherwise. Then \bar{y} is the sequence of moves for *Noselector* in a run of $G'(I)$ according to Σ_{a_i} . However, \bar{x} and \bar{y} are eventually intertwining, which by Lemma 2.3

implies that the comeager constant values for these two sequences are the same, giving a contradiction.

Let Z be the set of pairs $(x, a) \in I^+ \times ([X]^{<\omega} \setminus \{\emptyset\})$ such that *Noselector* has a winning strategy in the modified version of $G'(I)$ where *Selector* must play ω with his first move and *Noselector* must play x and declare a as the eventual constant value for \bar{x} . Let X' be the set of a for which there exists an x with $(x, a) \in Z$, and for each $a \in X'$, let J_a be the union of I with the set of x such that $(x, a) \in Z$. Then each J_a is an ideal, and the previous paragraph shows that every I -positive set contains an I -positive member of some J_a .

Another strategy-blending argument shows that the ideals J_a are pairwise I -orthogonal. Fix (x, a) and (x', a') in Z as witnessed (respectively) by strategies Δ and Δ' , and suppose that $x \cap x' \in I^+$.

Consider the sequences defined by

- $x_0 = x$
- $x'_0 = x'$
- $x_1 = \Delta(\langle \omega, x, x \cap x' \rangle)$
- $x'_1 = \Delta'(\langle \omega, x', x_1 \rangle)$
- $x_2 = \Delta(\langle \omega, x, x \cap x', x_1, x'_1 \rangle)$
- $x'_2 = \Delta'(\langle \omega, x', x_1, x'_1, x_2 \rangle)$
- $x_3 = \Delta(\langle \omega, x, x \cap x', x_1, x'_1, x_2, x'_2 \rangle)$
- $x'_3 = \Delta'(\langle \omega, x', x_1, x'_1, x_2, x'_2, x_3 \rangle)$

and so on. Then $\langle x_n : n \in \omega \rangle$ lists *Noselector's* moves for a run of $G'(I)$ according to Δ , $\langle x'_n : n \in \omega \rangle$ lists *Noselector's* moves for a run of $G'(I)$ according to Δ' , and the two sequences are eventually intertwining. Applying Lemma 2.3 again then completes the proof. \square

Given I, X, f , and $\mathcal{J} = \{J_a : a \in X'\}$ as in Theorem 3.2, f can be extended, using the map $x \mapsto C(I, \mathcal{J}, x)$, to a function on $\text{FCI}(I, \mathcal{J})$ which is invariant under finite changes and has codomain $[X]^{<\omega}$. Iterating this fact, possibly transfinitely, one must eventually add ω to the domain of the extended function. This gives the following result.

Theorem 3.3. *Suppose that $\text{AD}_{\mathbb{R}}$ holds. There is a definable operation which takes in a function of the form $f: (I \setminus \text{Fin}) \rightarrow ([X]^{<\omega} \setminus \{\emptyset\})$, where I is a tall ideal on ω containing Fin , X is a countable set and f is invariant under finite changes, and returns a function $f': (\mathcal{P}(\omega) \setminus \text{Fin}) \rightarrow ([X]^{<\omega} \setminus \{\emptyset\})$, also invariant under finite changes, such that $f'(x) = f(x)$ for all $x \in I \setminus \text{Fin}$.*

The following corollary was our motivation for proving this theorem.

Corollary 3.4. *Suppose that $\text{AD}_{\mathbb{R}}$ holds. Let Y be a set, and let G be a function with domain Y such that $G(y)$ is countable for all $y \in Y$. Suppose that there exists a $\mathcal{P}(\omega)/\text{Fin}$ -name for a function on Y which picks a nonempty finite subset of $G(y)$ for each $y \in Y$. Then there exists already a function on Y which picks a nonempty finite subset of $G(y)$ for each $y \in Y$.*

Proof. We show how to pick a finite subset of $G(y)$ for each $y \in Y$. Fix $y \in Y$. For each nonempty finite $a \subseteq G(y)$, let J_a be the union of Fin with the set of infinite $x \subseteq \omega$ forcing the value of τ at y to be a . Let \mathcal{J} be the collection of nonempty sets of the form J_a . Then \mathcal{J} is a collection of pairwise orthogonal ideals, and $\text{FCI}(\text{Fin}, \mathcal{J})$

is a tall ideal. Define

$$f: (\text{FCI}(\text{Fin}, \mathcal{J}) \setminus \text{Fin}) \rightarrow ([G(y)]^{<\omega} \setminus \{\emptyset\})$$

by setting $f(x) = \bigcup \{a \mid J_a \in C(I, \mathcal{J}, x)\}$. Then apply Theorem 3.3 with f , and let $f'(\omega)$ be the chosen set. \square

As a consequence, forcing with $\mathcal{P}(\omega)/\text{Fin}$ over a model of $\text{AD}_{\mathbb{R}}$ does not add an E_0 -selector. Furthermore, for any pair of equivalence relations E, F in a model M of $\text{AD}_{\mathbb{R}}$ for which every E -class is made up of countably many F -classes, if in a $\mathcal{P}(\omega)/\text{Fin}$ extension of M there is a function which picks a finite set of F -subclasses for each E -class, then such a function exists in M .

The argument for Corollary 3.4 also shows the following.

Corollary 3.5. *Suppose that $\text{AD}_{\mathbb{R}}$ holds. Then there is a function which takes in a countable family of orthogonal ideals whose union is dense in $\mathcal{P}(\omega)/\text{Fin}$ and returns a finite subfamily.*

Prikry [5] proved from $\text{AD}_{\mathbb{R}}$ that every set of reals is Ramsey. By work of Martin, Steel and Woodin this assumption was improved to AD^+ (see [1]). The following question asks for an analogous improvement for Theorem 3.2.

3.6 Question. Can the assumption of $\text{AD}_{\mathbb{R}}$ in Theorem 3.2 be replaced by AD^+ ?

Another natural question is the following.

3.7 Question. Are there uncountable sets X for which Theorem 3.3 holds?

4. EXAMPLES

By convention, Θ denotes the least ordinal which is not a surjective image of $\mathcal{P}(\omega)$. The iterated application of Theorem 3.2 in the proof of Theorem 3.3 clearly cannot run for Θ steps. Let us say that a function f as in Theorem 3.2 has *rank* α if it takes exactly α many iterations of Theorem 3.2 to add ω to its domain. We show in this section that, assuming $\text{AD}_{\mathbb{R}}$, there exist functions of rank α for cofinally many $\alpha < \Theta$. Our first example below illustrates a function of rank 1. The second shows how to combine a collection of functions, indexed by an almost disjoint family, to produce a function whose rank is greater than all the members of the collection.

4.1 Example. Suppose that $\{a_i : 1 \leq i < \omega\}$ is a partition of ω into infinite sets. For each positive $i \in \omega$, let J_i be the set of $x \subseteq \omega$ for which $x \setminus a_i$ is finite. Let J_0 be the set of $x \subseteq \omega$ having finite intersection with each a_i , and let $\mathcal{J} = \{J_i : i \in \omega\}$. Define $f: (\text{FCI}(\text{Fin}, \mathcal{J}) \setminus \text{Fin}) \rightarrow ([\mathcal{J}]^{<\omega} \setminus \{\emptyset\})$ by letting $f(x) = C(\text{Fin}, \mathcal{J}, x)$. Then every position (in particular, ω) in the game from the proof of Theorem 3.2 is a winning position for *Noselector* with value $\{J_0\}$, so f extends to all of $\mathcal{P}(\omega) \setminus \text{Fin}$ in one step.

4.2 Example. Suppose that we are given a countable set X (with $\emptyset \notin X$), a nonmaximal almost disjoint family \mathcal{A} of infinite subsets of ω (so distinct members of \mathcal{A} have finite intersection), and ideals I_a (properly containing Fin) and functions $f_a: (I_a \setminus \text{Fin}) \rightarrow ([X]^{<\omega} \setminus \{\emptyset\})$ ($a \in \mathcal{A}$) as in Theorem 3.2. Suppose that α_a ($a \in \mathcal{A}$)

are ordinals such that each f_a takes α_a many applications of Theorem 3.2 to add ω to its domain. Let J_\emptyset be \mathcal{A}^\perp , the set of $x \subseteq \omega$ having finite intersection with each $a \in \mathcal{A}$. For each $a \in \mathcal{A}$, let $\pi_a: \omega \rightarrow a$ be a bijection. For each $a \in \mathcal{A}$, let J_a be the ideal generated by $\text{Fin} \cup \{\pi_a[x] : x \in I_a\}$, and define $f'_a: (J_a \setminus \text{Fin}) \rightarrow ([X]^{<\omega} \setminus \{\emptyset\})$ by setting $f'_a(x) = f_a(\pi_a^{-1}[x \cap a])$. Let $\mathcal{J} = \{J_\emptyset\} \cup \{J_a : a \in \mathcal{A}\}$, and define $f: (\text{FCI}(\text{Fin}, \mathcal{J}) \setminus \text{Fin}) \rightarrow ([\{\emptyset\} \cup X]^{<\omega} \setminus \{\emptyset\})$ by letting

- $\emptyset \in f(x)$ if and only if $J_\emptyset \in C(\text{Fin}, \mathcal{J}, x)$,
- for all $z \in X$, $z \in f(x)$ if and only if there exists $a \in \mathcal{A}$ such that $J_a \in C(\text{Fin}, \mathcal{J}, x)$ and $z \in f'_a(x \cap a)$.

The iterated application of Theorem 3.2 for f adds subsets of each $a \in \mathcal{A}$ exactly when the corresponding process for f_a adds the π_a -preimages of these sets. It follows that it takes α_a many iterations of Theorem 3.2 to add each $a \in \mathcal{A}$ to the extended domain of f , and more than $\sup\{\alpha_a : a \in \mathcal{A}\}$ many iterations to add ω .

Fixing a bijection $g: \text{Fin} \rightarrow \omega$, the function which sends $a \in \mathcal{P}(\omega)$ to

$$\{g(a \cap n) : n \in \omega\}$$

is Borel and sends distinct infinite sets to infinite sets with finite intersection. It follows then (under ZF), that for each $\alpha < \Theta$ there is a surjection $\rho: \mathcal{A} \rightarrow \alpha$, for some almost disjoint family \mathcal{A} . Starting with a function as in Example 4.1, one can iteratively apply the construction from Example 4.2, filling in functions of rank β for the indices in $\rho^{-1}[\{\beta\}]$, to produce a function of rank at least α .

5. COUNTEREXAMPLES

The Axiom of Real Determinacy implies that every set of reals has the property of Baire, and this precludes a number of variations of the result proved in this paper. In this section we show that “finite subset of $G(y)$ ” cannot be replaced with “element of $[G(y)]^{\leq n} \setminus \{\emptyset\}$ ” in Corollary 3.4, for any positive $n \in \omega$, even if τ is a name for a function picking a single element from each $G(y)$. Similarly, “finite subfamily” cannot be replaced with “nonempty subfamily of size at most n ” in Corollary 3.5.

5.1 Example. Suppose that every set of reals has the property of Baire. Then there is no function $F: \mathcal{P}(\omega) \rightarrow 2$ with the property that for all subsets x, y of ω , $F(x) \neq F(y)$ if $x \Delta (\omega \setminus y)$ is finite (i.e., no function which chooses between complementary pairs of E_0 -degrees). This follows from the fact that F would be invariant under finite changes, so one of $F^{-1}[\{0\}]$ and $F^{-1}[\{1\}]$ would have to be comeager, but any comeager subset of $\mathcal{P}(\omega)$ contains a pair of complements. A similar argument shows (under the same assumption) that, for any $n \in \omega$, there is no function which takes in a collection $\{[a_i]_{E_0} : i < n\}$, for $\{a_i : i < n\}$ a partition of ω , and returns a nonempty proper subset. If there were such a function, there would be $j, k < n$ such that for comeagerly many partitions $\langle a_i : i < n \rangle$ of ω , $[a_j]_{E_0}$ would be in the chosen subset, and $[a_k]_{E_0}$ would not. However, comeagerly many members of this set would have the property that switching the values of a_j and a_k would give another member of the set, giving a contradiction.

To see this example in terms of Corollary 3.3, notice that a nonprincipal ultrafilter on ω induces a choice of one element from each set of the form

$$\{[a_i]_{E_0} : i \leq n\},$$

for $\{a_i : i \leq n\}$ a partition of ω into infinite sets. The previous paragraph shows that there is no function in a model of $\text{AD}_{\mathbb{R}}$ picking a proper subfamily from each such family.

The previous example does not give a counterexample corresponding to Theorem 3.3, as the corresponding ideal $\text{FCI}(\text{Fin}, \mathcal{J})$ would be all of $\mathcal{P}(\omega)$. We remedy this by showing how to replace each set of the form $[a]_{E_0}$ with an infinite maximal antichain in $\mathcal{P}(a)/E_0$. Once we have this, we can replace each set $\{[a_i]_{E_0} : i \leq n\}$ as above with the union of the corresponding antichains. Then as before, no function in the ground model can pick a subfamily of size at most n from each such union. The corresponding function f as in Theorem 3.3 has the form $x \mapsto C(\text{Fin}, \mathcal{J}, x)$ for $x \in \text{FCI}(\text{Fin}, \mathcal{J})$ and \mathcal{J} the collection of ideals derived from this antichain.

The only missing ingredient is a method for taking a set of the form $[a]_{E_0}$, for $a \subseteq \omega$ infinite, and returning a set of the form $[b]_{E_0}$, for some infinite $b \subseteq a$ for which $a \setminus b$ is infinite. The following observation does what we need.

Lemma 5.2. *There is a Borel function g that assigns each to infinite $a \subseteq \omega$ an infinite $g(a) \subseteq a$ for which $a \setminus g(a)$ is infinite, in such a way that $a \triangle b \in \text{Fin}$ implies that $g(a) \triangle g(b) \in \text{Fin}$.*

Proof. For each $n \in \omega$, let F_n be the equivalence relation on $\mathcal{P}(\omega)$ defined by $a F_n b \Leftrightarrow (a \setminus n) = (b \setminus n)$. Then define functions S_n ($n \in \omega$) on $\mathcal{P}(\omega)$ by

- $S_0(a)$ is the second element of a ,
- $S_{n+1}(a)$ is the second element of $\bigcap [a]_{F_{n+1}} \setminus (\sup\{S_n(b) + 1 : b \in [a]_{F_{n+1}}\})$.

Then for each infinite $a \subseteq \omega$ and each $n \in \omega$, $S_n(a) \geq 2n + 1$, and $\bigcap [a]_{F_{n+1}}$ can equivalently be replaced with a in the definition of $S_{n+1}(a)$. It follows that the function $g(a) = \{S_n(a) : n \in \omega\}$ is as desired. \square

REFERENCES

- [1] A. Caicedo, R. Ketchersid, *A trichotomy theorem in natural models of AD^+* , in: **Set Theory and Its Applications**, Contemporary Mathematics, vol. 533, Amer. Math. Soc., Providence, RI, 2011, 227-258
- [2] C.A. Di Prisco, S. Todorcevic, *Souslin partitions of products of finite sets*, Advances in Math 176 (2003), 145-173
- [3] P. Larson, D. Raghavan, *Real games and strategically selective coideals*, in preparation
- [4] A. Mathias, *Happy families*, Annals of Mathematical Logic 12 (1977), 59-111
- [5] K. Prikry, *Determinateness and partitions*, Proc. Amer. Math. Soc. 54 (1976), 303-306
- [6] S. Shelah, J. Zapletal, *Ramsey theorems for product of finite sets with submeasures*, Combinatorica 31 (2011) 225-244