

Majority decisions when abstention is possible

Paul Larson^{a,1}, Nick Matteo^b, Saharon Shelah^{c,2,3}

^a*Department of Mathematics, Miami University, Oxford, OH 45056, USA*

^b*Department of Mathematics, Miami University, Oxford, OH 45056, USA*

^c*The Hebrew University of Jerusalem, Einstein Institute of Mathematics*

Edmond J. Safra Campus, Givat Ram, Jerusalem 91904, Israel

Department of Mathematics, Hill Center-Busch Campus, Rutgers, The State University of New Jersey, 110 Frelinghuysen Road, Piscataway, NJ 08854, USA

Abstract

Suppose that we are given a family of choice functions on pairs from a given finite set. The set is considered as a set of alternatives (say candidates for an office) and the functions as potential “voters.” The question is, what choice functions agree, on every pair, with the majority of some finite subfamily of the voters? For the problem as stated, a complete characterization was given in Shelah [1], but here we allow voters to abstain. Aside from the trivial case, the possible families of (partial) choice functions break into three cases in terms of the functions that can be generated by majority decision. In one of these, cycles along the lines of Condorcet’s paradox are avoided. In another, all partial choice functions can be represented.

Keywords: choice function, majority decision, Condorcet’s paradox, tournament

2000 MSC: Primary 91B14, Secondary 05C20

1. Introduction

Condorcet’s “paradox” demonstrates that given three candidates A, B, and C, majority rule may result in the society preferring A to B, B to C, and C to A [2]. McGarvey [3] proved a far-reaching extension of Condorcet’s paradox: For every asymmetric relation R (so $a R b$ implies $b \not R a$) on a set X of \mathbf{n} candidates, there are m linear order relations on X : R_1, \dots, R_m , with R as their strict simple majority relation. That is, for every $a, b \in X$,

$$a R b \iff |\{i : a R_i b\}| > \frac{m}{2}.$$

Email addresses: laronpb@muohio.edu (Paul Larson), matteo.n@neu.edu (Nick Matteo)

¹The research of the first author is supported by NSF grant DMS-0801009.

²Partially supported by the United States-Israel Binational Science Foundation.

³This is publication number 965 in the third author’s publication list.

In other words, given any set of choices from pairs from a set of \mathbf{n} candidates, there is a population of m voters, all with simple linear-order preferences among the candidates, who will yield the given outcome for each pair in a majority-rule election between them.

McGarvey's proof gave $m = \mathbf{n}(\mathbf{n} - 1)$. Stearns [4] found a construction with $m = \mathbf{n}$ and noticed that a simple counting argument implies that m must be at least $\frac{\mathbf{n}}{\log \mathbf{n}}$. Erdős and Moser [5] were able to give a construction with $m = O\left(\frac{\mathbf{n}}{\log \mathbf{n}}\right)$. Mala [6] and Alon [7] showed that there is a positive constant c_1 such that, given any asymmetric relation R , there is some m and linear orders R_1, \dots, R_m with

$$a R b \iff |\{i : a R_i b\}| > \frac{m}{2} + c_1 \sqrt{m}$$

and that this is not true for any c_2 greater than c_1 .

Gil Kalai asked to what extent the assertion of McGarvey's theorem holds if we replace the linear orders by an arbitrary isomorphism class of choice functions on pairs of elements. Namely, when can we guarantee that every asymmetric relation R on X could result from a finite population of voters, each using a given kind of asymmetric relation on X ?

Informally, one may think of the situation we study here as follows. Imagine that we have \mathbf{n} candidates running for office, and that each voter votes in a two-step process as follows. First, the voter chooses one element from a set of ballot types, which are partial tournaments (or asymmetric relations) with \mathbf{n} vertices. Then, having chosen a ballot type, the voter labels the corresponding vertices with the names of the candidates, in such a way that an edge from candidate A to candidate B indicates a preference for A over B (of course, not all ballot types allow one to express one's preferences, but for the purposes of this illustration we assume that each voter does the best he or she can with the available options). The outcome of the election is the partial tournament with a node for each candidate and an edge from A to B if more voters had edges from A to B on their ballots than had the reverse. The question we address here is, what sets of ballot types can generate which partial tournaments via this process, assuming any number of voters? As we shall see, the nontrivial sets of ballot types (or, kinds of symmetric relations, as above) fall into three classes (see Main Claim 3.1 and the last paragraph of this introduction). Among these are the partisan case, in which voters are offered the chance to separate out a preferred subset of the candidates (which is how elections are often run, with the preferred set having size 1). We will show that the partisan case is the only one that does not lead to paradoxical results (i.e., directed cycles). We will also characterize the sets of ballot types which allow every partial tournament to be generated.

More formally, we use the following terminology. Let $\binom{X}{k}$ denote the family of subsets of X with k elements:

$$\binom{X}{k} = \{Y \subseteq X : |Y| = k\}.$$

Either an individual voter's preferences among candidates, or the outcomes that would result from each two-candidate election, may be represented as any of:

- An asymmetric relation R where $a R b$ if and only if a beats b ; it is possible that $a \not R b$, $b \not R a$, and $a \neq b$.
- A choice function defined on some subfamily of $\binom{X}{2}$, choosing the winner in each pair. Such a choice function is called “full” if its domain is all of $\binom{X}{2}$, and “partial” otherwise.
- An oriented graph, i.e. a directed graph with nodes X and edges $a \rightarrow b$ when a beats b .

We shall treat these representations as largely interchangeable throughout this paper. Total asymmetric relations, full choice functions, and tournaments (complete oriented graphs) all correspond to the case of no abstaining. Using notation from Shelah [1], we will let $\text{Tor}(c)$ denote the oriented graph (i.e., partial tournament) associated with a choice function c . For any set X , $c \mapsto \text{Tor}(c)$ is a bijection of full choice functions onto tournaments on X , and a bijection of all choice functions onto oriented graphs on X .

For the rest of this paper, we fix a finite set X . We let \mathbf{n} (as opposed to n , which is used as a variable) denote the size of X , and assume that $\mathbf{n} \geq 3$. We let \mathfrak{C} denote the set of choice functions on pairs from X , that is, the set of functions $c: Y \rightarrow X$ where $Y \subseteq \binom{X}{2}$ and $c\{x, y\} \in \{x, y\}$ for all $\{x, y\} \in Y$. When $c\{x, y\}$ is not defined it is interpreted as abstention or having no preference.

- Definition 1.1.** (a) $\text{Per}(X)$ is the set of permutations of X .
(b) A choice function d is a *permutation* of a choice function c if there is $\sigma \in \text{Per}(X)$ such that

$$d\{\sigma(x), \sigma(y)\} = \sigma(x) \iff c\{x, y\} = x;$$

we write $d = c^\sigma$.

- (c) A set of choice functions $\mathcal{C} \subseteq \mathfrak{C}$ is *symmetric* if it is closed under permutations of X . So for each $\sigma \in \text{Per}(X)$, if $c \in \mathcal{C}$ then $c^\sigma \in \mathcal{C}$.

Note that a choice function d is a permutation of a choice function c if and only if $\text{Tor}(c)$ and $\text{Tor}(d)$ are isomorphic directed graphs. Symmetric sets of choice functions are what was meant above by “kind of asymmetric relation.”

The main result of Shelah [1] pertained to full choice functions for the voters. It was shown that an arbitrary choice function d could result from a symmetric set \mathcal{C} consisting of full choice functions (i.e. for each d there is a finite set $\{c_1, \dots, c_m\} \subseteq \mathcal{C}$ such that $d\{x, y\} = x \iff |\{i : c_i\{x, y\} = x\}| > \frac{m}{2}$) if and only if for some $c \in \mathcal{C}$ and $x \in X$,

$$|\{y : c\{x, y\} = y\}| \neq \frac{\mathbf{n} - 1}{2}.$$

We shall call this condition “imbalance.”

Definition 1.2. For a choice function c ,

- (a) Let $\text{dom}(c)$ denote the domain of c .
- (b) for any pair (x, y) in X^2 , the *weight* of x over y for c is

$$W_y^x(c) = \begin{cases} 1 & \text{if } c\{x, y\} = x \\ 0 & \text{if } \{x, y\} \notin \text{dom } c \\ -1 & \text{if } c\{x, y\} = y \end{cases}$$

- (c) c is *balanced* if for all $x \in X$

$$\sum_{y \in X} W_y^x(c) = 0.$$

That is, $|\{y : c\{x, y\} = x\}| = |\{y : c\{x, y\} = y\}|$, for every x in X .

- (d) c is *unbalanced* if c is not balanced.
- (e) c is *pseudo-balanced* if every edge of $\text{Tor}(c)$ belongs to a directed cycle.

Gil Kalai further asked whether the number m can be given bounds in terms of \mathbf{n} , and what is the result of demanding a “non-trivial majority,” e.g. 51%. We shall consider loose bounds while addressing the general case, when voters are permitted to abstain.

To determine what symmetric sets of partial choice functions could produce an arbitrary outcome, we shall characterize the set of all possible outcomes of a symmetric set of choice functions, the “majority closure” $\text{maj-cl}(\mathcal{C})$. The symmetric set \mathcal{C} satisfies this extension to McGarvey’s theorem if and only if $\text{maj-cl}(\mathcal{C}) = \mathfrak{C}$.

Definition 1.3. For $\mathcal{C} \subseteq \mathfrak{C}$, let $\text{maj-cl}(\mathcal{C})$ be the set of $d \in \mathfrak{C}$ such that, for some set of weights $\{r_c \in [0, 1]_{\mathbb{Q}} : c \in \mathcal{C}\}$ with $\sum_{c \in \mathcal{C}} r_c = 1$,

$$d\{x, y\} = x \iff \sum_{c \in \mathcal{C}} W_y^x(c) r_c > 0.$$

Here and throughout this paper, $[0, 1]_{\mathbb{Q}}$ refers to the rationals in the unit interval of the real line. Note that we cannot assume that $\text{maj-cl}(\text{maj-cl}(\mathcal{C})) = \text{maj-cl}(\mathcal{C})$, and in fact we shall see this is not true. We must show that each function d in the majority closure, thus defined, can in fact be the outcome of a finite collection of voters using choice functions in \mathcal{C} ; this is Claim 2.6.

Shelah [1] gave a characterization of $\text{maj-cl}(\mathcal{C})$ for nonempty symmetric sets of full choice functions; there are just two cases. If some $c \in \mathcal{C}$ is unbalanced, then $\text{maj-cl}(\mathcal{C}) = \mathfrak{C}$; if every $c \in \mathcal{C}$ is balanced, then $\text{maj-cl}(\mathcal{C})$ is the set of all pseudo-balanced functions. The situation with possible abstention is more complicated; there are three nontrivial cases, see Main Claim 3.1. Unlike in Shelah [1], it is possible to have nontrivial \mathcal{C} for which $\text{maj-cl}(\mathcal{C})$ contains no directed cycles; this happens if and only if \mathcal{C} is partisan (see Definitions 2.3 and 2.4).

2. Basic Definitions and Facts

In addition to the representations as relations, choice functions, or oriented graphs discussed above, we shall also find it convenient to map each choice function c to a sequence in $[-1, 1]_{\mathbb{Q}}^{X^2}$ indexed by X^2 , the “probability sequence”

$$\text{pr}(c) = \langle W_y^x(c) : (x, y) \in X^2 \rangle.$$

Let $\text{pr}(\mathcal{C})$ be $\{\text{pr}(c) : c \in \mathcal{C}\}$, and $\text{pr}(X)$ denote the set of all sequences \bar{t} in $[-1, 1]_{\mathbb{Q}}^{X^2}$ such that $t_{x,y} = -t_{y,x}$; $\text{pr}(X)$ contains $\text{pr}(\mathcal{C})$.

Definition 2.1. For a probability sequence $\bar{t} \in \text{pr}(X)$,

- (a) \bar{t} is *balanced* if for each $x \in X$, $\sum_{y \in X} t_{x,y} = 0$.
- (b) $\text{maj}(\bar{t})$ is the $c \in \mathcal{C}$ such that $c\{x, y\} = x \iff t_{x,y} > 0$.

Note that maj and pr are mutually inverse functions from \mathcal{C} to $\text{pr}(\mathcal{C})$; $c = \text{maj}(\text{pr}(c))$ and $\bar{t} = \text{pr}(\text{maj}(\bar{t}))$ for all $c \in \mathcal{C}$ and $\bar{t} \in \text{pr}(\mathcal{C})$. The function maj is also defined on the much larger set $\text{pr}(X)$, which it maps onto \mathcal{C} . The reason for reusing the name “balanced” is clear:

Claim 2.2. *If $c \in \mathcal{C}$ is a balanced choice function, then $\text{pr}(c)$ is a balanced probability sequence. If $\bar{t} \in \text{pr}(\mathcal{C})$ is a balanced probability sequence, then $\text{maj}(\bar{t})$ is a balanced choice function.*

Proof. Suppose $c \in \mathcal{C}$ is balanced. For every $x \in X$, $\sum_{y \in X} W_y^x(c) = 0$. The (x, y) -term of $\text{pr}(c)$ is $W_y^x(c)$, so $\sum_{y \in X} \text{pr}(c)_{x,y} = 0$, i.e. $\text{pr}(c)$ is balanced.

Now suppose $\bar{t} \in \text{pr}(\mathcal{C})$ is balanced. Then $\bar{t} = \text{pr}(c)$ for some $c \in \mathcal{C}$. Since $t_{x,y} = W_y^x(c)$,

$$\sum_{y \in X} W_y^x(c) = \sum_{y \in X} t_{x,y} = 0$$

and c is balanced. □

However, it is not the case that $\text{maj}(\bar{t})$ is balanced for every balanced $\bar{t} \in \text{pr}(X)$. For instance, choose $x, z \in X$ and define a sequence \bar{t} which has $t_{z,x} = 1$, $t_{x,z} = -1$; for all $y \notin \{x, z\}$

$$\begin{aligned} t_{x,y} &= t_{y,z} = \frac{1}{\mathbf{n} - 2}, \\ t_{y,x} &= t_{z,y} = \frac{-1}{\mathbf{n} - 2}; \end{aligned}$$

and for all other pairs (u, v) , $t_{u,v} = 0$. One may check that \bar{t} is balanced, yet $\text{maj}(\bar{t})$ has $\sum_{y \in X} W_y^x(\text{maj}(\bar{t})) = \mathbf{n} - 3 > 0$ for any X with at least 4 elements.

Definition 2.3. For a choice function $c \in \mathcal{C}$,

- (a) c is *partisan* if there is nonempty $W \subsetneq X$ such that

$$c\{x, y\} = x \iff x \in W \text{ and } y \notin W.$$

- (b) c is *tiered* if there is a partition $\{X_1, X_2, \dots, X_k\}$ of X such that $c\{x, y\} = x$ if and only if $x \in X_i, y \in X_j$, and $i > j$. We say c is k -tiered where the partition has k sets. (So a partisan function is 2-tiered.)
- (c) c is *chaotic* if it is both unbalanced and nonpartisan.

Definition 2.4. For a subset $\mathcal{C} \subseteq \mathfrak{C}$,

- (a) \mathcal{C} is *trivial* if $\mathcal{C} = \emptyset$ or $\mathcal{C} = \{c\}$ where $\text{dom } c = \emptyset$, i.e. c makes no decisions.
- (b) \mathcal{C} is *balanced* if every $c \in \mathcal{C}$ is balanced.
- (c) \mathcal{C} is *partisan* if every $c \in \mathcal{C}$ is partisan.
- (d) \mathcal{C} is *chaotic* if there is some chaotic $c \in \mathcal{C}$.
- (e) $\text{pr-cl}(\mathcal{C})$ is the convex hull of $\text{pr}(\mathcal{C})$, i.e.

$$\text{pr-cl}(\mathcal{C}) = \left\{ \sum_{i=1}^k r_i \bar{t}_i : k \in \mathbb{N}, r_i \in [0, 1]_{\mathbb{Q}}, \sum_{i=1}^k r_i = 1, \bar{t}_i \in \text{pr}(\mathcal{C}) \right\}.$$

We can now establish some straightforward results which allow us to describe the possible outcomes due to voters chosen from a given symmetric set more explicitly.

Claim 2.5. *If \mathcal{C} is a symmetric subset of \mathfrak{C} , then d is the strict simple majority outcome of some finite set $\{c_1, \dots, c_m\}$ chosen from \mathcal{C} if and only if*

$$d\{x, y\} = x \iff \sum_{i=1}^m W_y^x(c_i) > 0.$$

Proof. We have that d is the strict simple majority outcome if and only if

$$\begin{aligned} d\{x, y\} = x &\iff |\{i : c_i\{x, y\} = x\}| > |\{i : c_i\{x, y\} = y\}| \\ &\iff \sum_{i=1}^m W_y^x(c_i) > 0. \end{aligned}$$

□

Claim 2.6. *Suppose \mathcal{C} is a symmetric subset of \mathfrak{C} . Then $d \in \text{maj-cl}(\mathcal{C})$ if and only if d is a strict simple majority outcome of some $\{c_1, \dots, c_m\} \subseteq \mathcal{C}$.*

Proof. If $d \in \mathfrak{C}$ and there is a set $\{c_1, \dots, c_m\} \subseteq \mathcal{C}$ with d as their strict simple majority outcome, then for each $c \in \mathcal{C}$ let r_c be the number of times that c appears among the c_i , divided by m :

$$r_c = \frac{|\{i : c_i = c\}|}{m}.$$

Clearly $\forall c \in \mathcal{C} \ 0 \leq r_c \leq 1$ and $\sum_{c \in \mathcal{C}} r_c = 1$. Furthermore,

$$d\{x, y\} = x \iff \sum_{i=1}^m W_y^x(c_i) > 0 \iff \sum_{i=1}^m \frac{W_y^x(c_i)}{m} > 0 \iff \sum_{c \in \mathcal{C}} W_y^x(c) r_c > 0.$$

Conversely, suppose $d \in \text{maj-cl}(\mathcal{C})$. There are $r_c = \frac{a_c}{b_c}$ for each $c \in \mathcal{C}$ with $\sum r_c = 1$, and

$$d\{x, y\} = x \iff \sum_{c \in \mathcal{C}} W_y^x(c) r_c > 0.$$

Let $m = \text{lcm}\{b_c : c \in \mathcal{C}\}$. Now construct a finite set $\{c_1, \dots, c_m\}$ by taking $a_c \frac{m}{b_c}$ copies of each $c \in \mathcal{C}$. $\frac{m}{b_c}$ is an integer since $b_c \mid m$. Now for each $c \in \mathcal{C}$,

$$\sum_{c_i=c} W_y^x(c_i) = W_y^x(c) \cdot |\{i : c_i = c\}| = W_y^x(c) a_c \frac{m}{b_c} = W_y^x(c) r_c m.$$

So

$$\begin{aligned} \sum_{i=1}^m W_y^x(c_i) &= \sum_{c \in \mathcal{C}} \sum_{c_i=c} W_y^x(c_i) = \sum_{c \in \mathcal{C}} W_y^x(c) r_c m. \\ \sum_{c \in \mathcal{C}} W_y^x(c) r_c m > 0 &\iff \sum_{c \in \mathcal{C}} W_y^x(c) r_c > 0, \end{aligned}$$

so d is the simple majority relation of the c_i . □

Claim 2.7. For any symmetric $\mathcal{C} \subseteq \mathfrak{C}$, $\text{maj-cl}(\mathcal{C}) = \{\text{maj}(\bar{t}) : \bar{t} \in \text{pr-cl}(\mathcal{C})\}$.

Proof. Suppose $d \in \text{maj-cl}(\mathcal{C})$. Then there are r_c per $c \in \mathcal{C}$ with $\sum_{c \in \mathcal{C}} r_c = 1$ and

$$d\{x, y\} = x \iff \sum_{c \in \mathcal{C}} r_c W_y^x(c) > 0.$$

Let $\bar{t} = \sum_{c \in \mathcal{C}} r_c \text{pr}(c)$ be a probability sequence in $\text{pr-cl}(\mathcal{C})$. Then

$$t_{x,y} = \sum_{c \in \mathcal{C}} r_c \text{pr}(c)_{x,y} = \sum_{c \in \mathcal{C}} r_c W_y^x(c).$$

So

$$\text{maj}(\bar{t})\{x, y\} = x \iff \sum_{c \in \mathcal{C}} r_c W_y^x(c) > 0,$$

i.e. $d = \text{maj}(\bar{t})$.

Suppose $d = \text{maj}(\bar{t})$ for some $\bar{t} \in \text{pr-cl}(\mathcal{C})$. Then for some r_c with $\sum_{c \in \mathcal{C}} r_c = 1$, $d = \text{maj}(\sum_{c \in \mathcal{C}} r_c \text{pr}(c))$.

$$d\{x, y\} = x \iff \sum_{c \in \mathcal{C}} r_c \text{pr}(c)_{x,y} > 0 \iff \sum_{c \in \mathcal{C}} r_c W_y^x(c) > 0.$$

So $d \in \text{maj-cl}(\mathcal{C})$. □

3. A Characterization

Main Claim 3.1. *Given a symmetric $\mathcal{C} \subseteq \mathfrak{C}$,*

- (i) \mathcal{C} is trivial \iff $\text{maj-cl}(\mathcal{C})$ is trivial.
- (ii) \mathcal{C} is balanced but nontrivial \iff $\text{maj-cl}(\mathcal{C})$ is the set of all pseudo-balanced choice functions on X .
- (iii) \mathcal{C} is partisan and nontrivial \iff $\text{maj-cl}(\mathcal{C})$ is the set of all tiered choice functions on X .
- (iv) \mathcal{C} is not balanced and not partisan \iff $\text{maj-cl}(\mathcal{C}) = \mathfrak{C}$.

Proof. No nontrivial \mathcal{C} is both balanced and partisan, so the sets

$$\begin{aligned} & \{\mathcal{C} \subset \mathfrak{C} : \mathcal{C} \text{ is trivial}\}, \\ & \{\mathcal{C} \subset \mathfrak{C} : \mathcal{C} \text{ is symmetric, nontrivial, and balanced}\}, \\ & \{\mathcal{C} \subset \mathfrak{C} : \mathcal{C} \text{ is symmetric, nontrivial, and partisan}\}, \text{ and} \\ & \{\mathcal{C} \subseteq \mathfrak{C} : \mathcal{C} \text{ is symmetric, not balanced, and not partisan}\} \end{aligned}$$

partition the family of all symmetric subsets of \mathfrak{C} . Thus proving each forward implication in the claim will give the reverse implications.

i. If no-one in \mathcal{C} makes any choices, no combination can have a majority choice.

ii. This is the content of Section 4, below.

iii. It is shown in Inada [8] (using different terminology) that if \mathcal{C} is partisan and nontrivial then every element of $\text{maj-cl}(\mathcal{C})$ is a tiered choice function. We give an alternate proof. Suppose that \mathcal{C} is partisan and $d \in \text{maj-cl}(\mathcal{C})$. Then there is a finite set $\{c_1, \dots, c_m\} \subseteq \mathcal{C}$ with d as the strict simple majority outcome. For each $x \in X$, let k_x be the number of c_i such that x is in the “winning” partite set. For any $(x, y) \in X^2$, let k be the number of c_i where x and y are both in the winning subset; the number of c_i with $c_i\{x, y\} = x$ is $k_x - k$, and the number with $c_i\{x, y\} = y$ is $k_y - k$, so

$$d\{x, y\} = x \iff k_x - k > k_y - k \iff k_x > k_y.$$

Partition X into subsets for each value of k_x ,

$$X = \bigcup \{\{x \in X : k_x = k\} : k \in \{k_x : x \in X\}\}.$$

These subsets form the tiers, so d is a tiered function.

For the reverse inclusion, suppose d is an arbitrary tiered function with tiers $\{X_1, \dots, X_k\}$. Let $c \in \mathcal{C}$. For each $x \in X$, let $\Gamma_x \subset \text{Per}(X)$ be all permutations holding x fixed and c_x be a choice function in \mathcal{C} with x in its winning subset. (If y is in the winning subset of c , $c^{(x,y)}$ will suffice for c_x .) Now construct $\{c_1, \dots, c_m\}$ by taking, for $1 \leq i \leq k$, each $x \in X_i$, and each $\sigma \in \Gamma_x$, i copies of c_x^σ . Suppose $x \in X_i$, $y \in X_j$, and $i > j$ (so $d\{x, y\} = x$). Then for any $z \in X_i \setminus \{x\}$, any $z \in X_j \setminus \{y\}$, and any z in other tiers, x and y are chosen by equal numbers of $\{c_z^\sigma : \sigma \in \Gamma_z\}$. So they are chosen by an equal number of

$\{c_1, \dots, c_m\}$ except among the functions c_x^σ and c_y^σ ; there are i such functions c_x^σ for each $\sigma \in \Gamma_x$, and j such functions c_y^σ for each $\sigma \in \Gamma_y$. The number of permutations fixing x and the number fixing y are the same, $|\Gamma_x| = |\Gamma_y|$. The number of permutations fixing x and putting y in the losing subset and the number fixing y and putting x in the losing subset is the same, l . There are il functions which choose x over y , and jl which choose y over x ; $il > jl$ so the strict simple majority outcome of the c_i chooses x .

If $i = j$, then $il = jl$, so x and y tie in the majority outcome. Thus d is the strict simple majority outcome of $\{c_1, \dots, c_m\}$ and $d \in \text{maj-cl}(\mathcal{C})$.

iv. If \mathcal{C} is a chaotic set, the conclusion follows from Section 5, below.

Otherwise, \mathcal{C} is not balanced and not partisan, but not chaotic. In this case, \mathcal{C} contains unbalanced functions which are all partisan, and nonpartisan functions which are all balanced. By symmetry, \mathcal{C} contains a nontrivial balanced symmetric subset \mathcal{B} and a nontrivial partisan symmetric subset \mathcal{P} .

Let $d \in \mathcal{C}$. Suppose $b \rightarrow a$ is any edge of d . For each $z \in X \setminus \{a, b\}$, by Claim 4.5 there is a voter population T_z from \mathcal{B} such that a beats z , z beats b , b beats a , and no other pairs are decided; moreover, the same number of voters, say m , choose the winner in each pair, and no dissenting votes occur. In the combination $\bigcup_z T_z$, m voters choose a over each z , and each z over b . $m(\mathbf{n} - 2)$ voters choose b over a .

Let $c \in \mathcal{P}$ be a partisan function, and l be the size of the set of winning candidates under c . Let $A \subset \mathcal{P}$ be the set of permutations of c so that a is on the losing side, $B \subset \mathcal{P}$ be the permutations of c so that b is on the winning side, and $C = A \cap B$ —the permutations with b winning and a losing.

$$|A| = \binom{\mathbf{n} - 1}{l}, \quad |B| = \binom{\mathbf{n} - 1}{l - 1}, \quad |C| = \binom{\mathbf{n} - 2}{l - 1}.$$

Any candidates besides a or b are tied over all the voters of A , B , or C , since each is selected in the winning set an equal number of times. Furthermore, a does not beat b in any choice function in A , B , or C .

Now we seek to take appropriate numbers of copies of the sets A , B , and C so that, for some constant k , any $z \in X \setminus \{a, b\}$ defeats a by k votes, and loses to b by k votes. Say we have k_0 copies of A , k_1 copies of B , and k_2 copies of C in a population D . Now for each $z \in X \setminus \{a, b\}$,

$$\begin{aligned} |\{c \in D : c\{z, a\} = z\}| &= k_0 \binom{\mathbf{n} - 2}{l - 1} + k_1 \binom{\mathbf{n} - 3}{l - 2} + k_2 \binom{\mathbf{n} - 3}{l - 2}, \\ |\{c \in D : c\{z, a\} = a\}| &= k_1 \binom{\mathbf{n} - 3}{l - 2}, \\ |\{c \in D : c\{b, z\} = b\}| &= k_0 \binom{\mathbf{n} - 3}{l - 1} + k_1 \binom{\mathbf{n} - 2}{l - 1} + k_2 \binom{\mathbf{n} - 3}{l - 1}, \\ |\{c \in D : c\{b, z\} = z\}| &= k_0 \binom{\mathbf{n} - 3}{l - 1}. \end{aligned}$$

So we solve $k_0 \binom{\mathbf{n} - 2}{l - 1} + k_2 \binom{\mathbf{n} - 3}{l - 2} = k_1 \binom{\mathbf{n} - 2}{l - 1} + k_2 \binom{\mathbf{n} - 3}{l - 1}$, with coefficients not all

zero. There are solutions

$$\begin{aligned} k_0 &= \mathbf{n} - 2l, & k_1 &= 0, & k_2 &= \mathbf{n} - 2, & \text{if } l &\leq \frac{\mathbf{n}}{2}; \\ k_0 &= 0, & k_1 &= 2l - \mathbf{n}, & k_2 &= \mathbf{n} - 2, & \text{if } l &> \frac{\mathbf{n}}{2}. \end{aligned}$$

Let $k = k_0 \binom{\mathbf{n}-2}{l-1} + k_2 \binom{\mathbf{n}-3}{l-2}$, the number of votes for b over z , or z over a , for any $z \in X \setminus \{a, b\}$. Now take the union of k copies of the population $\bigcup_z T_z$, mk_0 copies of A , mk_1 copies of B , and mk_2 copies of C for our voter population. b defeats any z mk times among the partisan functions, and z defeats b km times among the balanced functions, so they tie; similarly, a and z tie. Thus we are left with $b \rightarrow a$. The union of such populations for each edge in d yields d as the majority outcome.

Hence $\mathfrak{C} \subseteq \text{maj-cl}(\mathcal{C})$. \square

4. Balanced Choices

Definition 4.1. For a choice function $c \in \mathfrak{C}$,

- (a) c is *triangular* if for some $\{x, y, z\} \in \binom{X}{3}$, $c\{x, y\} = x$, $c\{y, z\} = y$, $c\{z, x\} = z$, and for any $\{u, v\} \not\subseteq \{x, y, z\}$, $\{u, v\} \notin \text{dom } c$. We write $c^{x,y,z}$ for such c .
- (b) c is *cyclic* if for some $\{x_1, \dots, x_k\} \in \binom{X}{k}$,
 - $c\{x_i, x_j\} = x_i$ if $j \equiv i + 1 \pmod k$.
 - No other pair $\{x, y\}$ is in $\text{dom } c$.

We write c^{x_1, \dots, x_k} for such c . (Triangular functions are a specific case of cyclic functions.)

- (c) Let pr^{bl} be the set of all balanced $\bar{t} \in \text{pr}(X)$.

Note that if \mathcal{C} is symmetric and $c^{x,y,z} \in \mathcal{C}$, then $c^{u,v,w} \in \mathcal{C}$ for any $\{u, v, w\} \in \binom{X}{3}$; similarly for cyclic functions.

Claim 4.2. *If a choice function $c = \text{maj}(\bar{t})$ for some $\bar{t} \in \text{pr}^{\text{bl}}$, then c is pseudo-balanced.*

Proof. Assume $c = \text{maj}(\bar{t})$ for some $\bar{t} \in \text{pr}^{\text{bl}}$. Let $x \rightarrow y$ be any edge of $\text{Tor}(c)$; then $t_{x,y} > 0$. Suppose $x \rightarrow y$ is in no directed cycle. Let Y be the set of $z \in X$ with a winning chain to x , i.e.

$$Y = \bigcup \left\{ \{z_1, \dots, z_k\} \in \binom{X \setminus \{y\}}{k} : k < \mathbf{n}, \forall i < k \ c\{z_i, z_{i+1}\} = z_i, z_k = x \right\}.$$

Let $\bar{Y} = X \setminus Y$. Note that $y \in \bar{Y}$ and $x \in Y$, so Y and \bar{Y} partition X into nonempty sets. Suppose $z \in Y$ and $v \in \bar{Y}$. Say z, z_1, \dots, z_k is a winning chain from z to x . If $t_{z,v} < 0$, then v, z, z_1, \dots, z_k forms a winning chain from v to x . If v is y , the chain forms a directed cycle with the edge $x \rightarrow y$ and we are done. If $v \neq y$, this contradicts that $v \notin Y$. So we assume that $t_{z,v} \geq 0$.

Now

$$\sum_{z \in Y, v \in \bar{Y}} t_{z,v} > 0,$$

since every $t_{z,v} \geq 0$, and $t_{x,y} > 0$ is among them. For each $u \in \bar{Y}$ let

$$\begin{aligned} r_u &= \sum_{z \in Y} t_{z,u}, \\ \bar{r}_u &= \sum_{v \in \bar{Y}} t_{v,u}. \end{aligned}$$

Since \bar{t} is balanced, $r_u + \bar{r}_u = 0$, so

$$\sum_{u \in \bar{Y}} (r_u + \bar{r}_u) = 0 = \sum_{u \in \bar{Y}} r_u + \sum_{u \in \bar{Y}} \bar{r}_u.$$

We have seen that the first summand is positive, so the second summand is negative. But it is zero because for each pair $(u, v) \in \bar{Y}^2$ we have $t_{u,v} + t_{v,u} = 0$. This is a contradiction; so $x \rightarrow y$ must be in some directed cycle. \square

Claim 4.3. pr^{bl} is a convex subset of $\text{pr}(X)$.

Proof. Suppose \bar{t} and \bar{s} are balanced probability sequences, and $a \in [0, 1]_{\mathbb{Q}}$. Then $\bar{u} = a\bar{t} + (1-a)\bar{s}$ has, for any $x \in X$,

$$\begin{aligned} \sum_{y \in X} u_{x,y} &= \sum_{y \in X} (at_{x,y} + (1-a)s_{x,y}) \\ &= a \sum_{y \in X} t_{x,y} + (1-a) \sum_{y \in X} s_{x,y} \\ &= a0 + (1-a)0 = 0. \end{aligned}$$

So \bar{u} is a balanced probability sequence. \square

Claim 4.4. If $\mathcal{C} \subset \mathfrak{C}$ is symmetric, balanced, and nontrivial, then every $d \in \text{maj-cl}(\mathcal{C})$ is pseudo-balanced.

Proof. By Claim 2.2, every $\bar{t} \in \text{pr}(\mathcal{C})$ is balanced. So by Claim 4.3, the convex hull $\text{pr-cl}(\mathcal{C})$ is contained in pr^{bl} . Any $d \in \text{maj-cl}(\mathcal{C})$ is $\text{maj}(\bar{t})$ for some $\bar{t} \in \text{pr-cl}(\mathcal{C})$, by Claim 2.7, hence is pseudo-balanced, by Claim 4.2. \square

For the reverse inclusion, that every pseudo-balanced function is in the majority closure, we shall consider the graph interpretation. First we shall show that every triangular function is in the majority closure of a balanced symmetric set.

Claim 4.5. If $\mathcal{C} \subset \mathfrak{C}$ is symmetric, balanced, and nontrivial, then for any $\{x, y, z\} \in \binom{X}{3}$, the triangular function $c^{x,y,z} \in \text{maj-cl}(\mathcal{C})$. Moreover, there is a voter population generating $c^{x,y,z}$ such that the same number of votes choose x over y , y over z , or z over x , and there are no opposing votes in any of these cases.

$y \quad z$

x

Let $\Gamma_{x,y}$,
 $1 \leq i \leq$
outcome

If u, v
 $c_i^\sigma\{u, v\} =$
not, then
are an eq
edges wh
where it c
no edges
only in

- c_1^σ ,
the
- c_2^σ ,
the

x_3

x_0

x_2

simple majority outcome. Let

$$l_1 = \sum_{i=1}^{m_1} W_{x_{k-1}}^{x_1}(c'_i).$$

Like l_0 , l_1 is a positive integer. Let $L = \text{lcm}(l_0, l_1)$, and take $\frac{L}{l_0}$ copies of each c_i , and $\frac{L}{l_1}$ copies of each c'_i , to make a finite set of voters $\{d_1, \dots, d_m\}$ where $m = \frac{L}{l_0}m_0 + \frac{L}{l_1}m_1$. Let d be their strict simple majority outcome. For any $\{u, v\} \in \binom{X}{2}$, if $\{u, v\} \not\subset \{x_1, \dots, x_k\}$ then u ties v among the c_i and the c'_i , so $\{u, v\} \notin \text{dom } d$. Any two points on the cycle (x_1, \dots, x_{k-1}) which are not adjacent are tied among the c_i and among the c'_i . Any point on the cycle besides x_1 or x_{k-1} is also tied with x_k among the c_i and among the c'_i . Any pair of consecutive points (x_i, x_j) on the cycle, besides (x_{k-1}, x_1) , are tied among the c'_i , but the majority of the c_i pick x_i . So $d\{x_i, x_j\} = x_i$. x_{k-1} and x_k tie among the c_i , but the majority of the c'_i pick x_{k-1} , so $d\{x_{k-1}, x_k\} = x_{k-1}$. Similarly $d\{x_k, x_1\} = x_k$. The only remaining pair to consider is $\{x_1, x_{k-1}\}$. $d\{x_1, x_{k-1}\}$ is defined if and only if

$$\sum_{i=1}^m W_{x_{k-1}}^{x_1}(d_i) \neq 0.$$

The left hand side is equal to

$$\begin{aligned} \frac{L}{l_0} \sum_{i=1}^{m_0} W_{x_{k-1}}^{x_1}(c_i) + \frac{L}{l_1} \sum_{i=1}^{m_1} W_{x_{k-1}}^{x_1}(c'_i) \\ = \frac{L}{l_0}(-l_0) + \frac{L}{l_1}(l_1) = 0. \end{aligned}$$

Thus $\{x_1, x_{k-1}\} \notin \text{dom } d$, and $d\{x_i, x_j\} = x_i$ if and only if $j \equiv i + 1 \pmod k$, so $d = c^{x_1, \dots, x_k} \in \text{maj-cl}(\mathcal{C})$.

By induction on k , any cyclic choice function on X is in $\text{maj-cl}(\mathcal{C})$. \square

Claim 4.7. *If $\mathcal{C} \subset \mathfrak{C}$ is symmetric, balanced, and nontrivial, then every pseudo-balanced choice function $d \in \mathfrak{C}$ is in $\text{maj-cl}(\mathcal{C})$.*

Proof. Suppose d is an arbitrary pseudo-balanced choice function. Every edge of $\text{Tor}(d)$ is on a directed cycle, so consider the decomposition of $\text{Tor}(d)$ into cycles $C_{x,y}$ for each edge $x \rightarrow y$ in $\text{Tor}(d)$. Let $d_{x,y}$ be the cyclic function corresponding to $C_{x,y}$ for each edge $x \rightarrow y$ in $\text{Tor}(d)$. By Claim 4.5, \mathcal{C} meets the requirements of Claim 4.6, so every cyclic function on X is in $\text{maj-cl}(\mathcal{C})$; in particular, each $d_{x,y} \in \text{maj-cl}(\mathcal{C})$. If we combine all the finite sets of voters which yielded each $d_{x,y}$, we get another finite set of voters from \mathcal{C} , since there are finitely many edges in $\text{Tor}(d)$.

For any $\{u, v\} \in \binom{X}{2} \setminus \text{dom } d$, the edge $u \rightarrow v$ is not in any cycle $C_{x,y}$. We have that $\sum W_v^u(c_i) = 0$ over the c_i yielding any $d_{x,y}$, so $\sum W_v^u(c_i) = 0$ over all our voters and u and v tie in their strict simple majority outcome.

For any $\{u, v\} \in \binom{X}{2}$ with $d\{u, v\} = u$, the edge $u \rightarrow v$ is in $\text{Tor}(d)$. Since each cycle $C_{x,y}$ has edges only from $\text{Tor}(d)$, each population of c_i yielding some $d_{x,y}$ has $\sum W_v^u(c_i) \geq 0$, and in particular the population yielding $d_{u,v}$ has $\sum W_v^u(c_i) > 0$, so over all our voters $\sum W_v^u(c_i) > 0$.

Thus the strict simple majority outcome is d , and $d \in \text{maj-cl}(\mathcal{C})$. \square

Combining Claims 4.4 and 4.7, we have that $\text{maj-cl}(\mathcal{C})$ is the set of all pseudo-balanced choice functions whenever \mathcal{C} is symmetric, balanced, and non-trivial.

5. Chaotic Choices

Definition 5.1. For a choice function $c \in \mathfrak{C}$,

(a) For $x \in X$, the *valence* of x with c is

$$\text{val}_c(x) = \sum_{y \in X} W_y^x(c).$$

(b) For $\ell \in \{-1, 0, 1\}$, let

$$V_\ell(c) = \{(\text{val}_c(x) - \ell, \text{val}_c(y) + \ell) : \{x, y\} \in \binom{X}{2}, W_y^x(c) = \ell\}.$$

These are the valence pairs, with winners on the right in V_{-1} , ties in V_0 , winners on the left in V_1 , and with 1 subtracted from the valence of the winner. In fact,

$$V_\ell(c) = \left\{ \sum_{z \in X \setminus \{x, y\}} (W_z^x(c), W_z^y(c)) : \{x, y\} \in \binom{X}{2}, W_y^x(c) = \ell \right\}.$$

(c) For a subset A of $\mathbb{Q} \times \mathbb{Q}$, let $\text{conv}(A)$ be the convex hull of A in $\mathbb{Q} \times \mathbb{Q}$.

(d) Let

$$V^*(c) = \{a\bar{v}_1 + (1-a)\bar{v}_0 : \bar{v}_1 \in \text{conv}(V_1(c)), \bar{v}_0 \in \text{conv}(V_0(c)), a \in (0, 1]_{\mathbb{Q}}\}.$$

These are the convex hulls of the “winning” valence pairs, together with ties, requiring some contribution from a non-tied pair.

Claim 5.2. For each $\ell \in \{-1, 0, 1\}$, $(k_0, k_1) \in V_\ell(c) \iff (k_1, k_0) \in V_{-\ell}(c)$.

Proof. A pair $(\text{val}_c(u) + 1, \text{val}_c(v) - 1)$ is in $V_{-1}(c)$ if and only if $W_v^u(c) = -1 \iff W_u^v(c) = 1$, so $(\text{val}_c(v) - 1, \text{val}_c(u) + 1) \in V_1(c)$.

A pair $(\text{val}_c(u), \text{val}_c(v))$ is in $V_0(c)$ if and only if $\{u, v\} = \{v, u\} \notin \text{dom } c$, so $(\text{val}_c(v), \text{val}_c(u)) \in V_0(c)$. \square

Claim 5.3. An unbalanced $c \in \mathfrak{C}$ is partisan if and only if

- $V_1(c)$ lies on a line parallel to the line $y = x$, and

- $V_0(c)$ is contained in the line $y = x$.

Proof. Suppose that $V_0(c)$ is all on $y = x$. Thus, candidates are only ever tied with others of the same valence. Suppose further that $V_1(c)$ is on a line $y = x - b$. Then whenever $c\{w, v\} = w$, $\text{val}_c(w) - \text{val}_c(v) = b + 2$ is constant; the valence of any winner is always $b + 2$ more than the defeated. Since c is unbalanced, there is such a pair $\{w, v\}$. If any candidate z has a valence different from that of w , it cannot be tied with w , so $\{w, z\} \in \text{dom } c$. Therefore the valences differ by $b + 2$, but if $\text{val}_c(z) \neq \text{val}_c(v)$, then z can neither tie nor be comparable to v , a contradiction. So every candidate has the valence of w or the valence of v . If two candidates with the same valence do not tie, then $b = -2$ and $\text{val}_c(v) = \text{val}_c(w)$. In this case, every candidate has the same valence, which would mean that c is balanced; a contradiction. Furthermore, every candidate with the high valence must defeat everyone of the low valence, since they cannot be tied. This is the definition of a partisan function.

Conversely, suppose that c is partisan. Then two elements are tied only if they are both in the winning subset, or both in the losing subset; in either case, they have the same valence, so $V_0(c)$ is contained in the line $y = x$. $V_1(c)$ is the single point $(\text{val}_c(w) - 1, \text{val}_c(v) + 1)$ for any w in the winning subset and v in the losing subset. Naturally, this point is contained in a line parallel to $y = x$. \square

Claim 5.4. *If $c \in \mathfrak{C}$ is unbalanced, then a point of $V^*(c)$ lies above the line $y = -x$, and a point of $V^*(c)$ lies below it.*

Proof. Let v_1 and v_2 be two candidates in X having the highest valences with c . Since c is unbalanced, $\text{val}_c(v_1) > 0$. The sum of the corresponding pair in any $V_\ell(c)$ is $\text{val}_c(v_1) + \text{val}_c(v_2)$. If this is less than or equal to 0, then $\text{val}_c(v_2) \leq -1$. Thus all other valences are at most -1 , so the average valence is strictly smaller than the average of $\text{val}_c(v_1)$ and $\text{val}_c(v_2)$, at most 0. This is a contradiction, since the average valence is always 0. Therefore, the sum of the valence pair for v_1 and v_2 in any $V_\ell(c)$ satisfies $y + x > 0$. If the pair is in $V_1(c)$, it is in $V^*(c)$. If it is in $V_{-1}(c)$, then its reflection in $V_1(c)$ (hence in $V^*(c)$) still satisfies $x + y > 0$. If it is in $V_0(c)$, then points arbitrarily close to the pair, along any line to a point of $V_1(c)$, are in $V^*(c)$. Some of these points are above $y = -x$.

Let u_1 and u_2 be two candidates in X having the smallest valences with c . By imbalance, $\text{val}_c(u_1) < 0$. Suppose $\text{val}_c(u_1) + \text{val}_c(u_2) \geq 0$. Then the average of these two valences is at least 0, and the average over all candidates is larger; it is strictly larger, since $\text{val}_c(v_1) > 0$ figures in the average. This is a contradiction, so the sum of the valence pair for u_1 and u_2 in any $V_\ell(c)$ satisfies $y + x < 0$. If this pair is in $V_{-1}(c)$, then its reflection in $V_1(c)$ still satisfies $x + y < 0$. Otherwise it is in $V^*(c)$, or arbitrarily close points are in $V^*(c)$. \square

To establish that every choice function is in the majority closure of a chaotic symmetric set, we shall first establish that some c therein satisfies a rather abstruse condition which we'll call "valence-imbalance."

\bar{p}_1

\bar{p}_0

$y = x$

for $n \geq 1$, approaching \bar{u} as $n \rightarrow \infty$. For each n , let \bar{v}_n be the intersection of the line through \bar{u}_n and \bar{w} with the line L . (If necessary, define the \bar{v}_n only for $n > j$, if the line from \bar{u}_j through \bar{w} is parallel to L ; at most one such j can exist.) As $n \rightarrow \infty$, \bar{v}_n approaches \bar{v} . So for some $N \in \mathbb{N}$ and all $n \geq N$, \bar{v}_n is in the interval (\bar{q}_0, \bar{q}_1) , an open neighborhood of \bar{v} in L . So

$$\bar{w} = r_{-1}\bar{u}_N + r_1\bar{v}_N,$$

for some r_{-1} and r_1 in $[0, 1]_{\mathbb{Q}}$ with $r_{-1} + r_1 = 1$.

If $r_{-1} = r_1 = \frac{1}{2}$, then $\bar{v}_N = \bar{w} + (\bar{w} - \bar{u}_N)$ is just as far from the line $y = x$ as $\bar{u}_N = \bar{w} - (\bar{w} - \bar{u}_N)$. But notice, either

- both the interval (\bar{u}, \bar{p}_1) on Λ and the interval (\bar{v}, \bar{q}_1) on L are farther from $y = x$ than \bar{u} is, while the intervals (\bar{p}_0, \bar{u}) and (\bar{q}_0, \bar{v}) are closer, or
- the intervals (\bar{u}, \bar{p}_1) and (\bar{v}, \bar{q}_1) are closer to $y = x$ than \bar{u} is, while the intervals (\bar{p}_0, \bar{u}) and (\bar{q}_0, \bar{v}) are farther.

Then \bar{u}_N is in the half-plane $y + x > 2a + b$, and \bar{w} is the sole intersection of the line $\{\bar{u}_N + t(\bar{w} - \bar{u}_N) : t \in \mathbb{Q}\}$ with the line $y + x = 2a + b$. Thus \bar{v}_N is in the half-plane $y + x < 2a + b$, hence in the interval (\bar{q}_0, \bar{v}) of L . But then the distance from $y = x$ to \bar{u} is strictly between the distances from $y = x$ to \bar{v}_N and to \bar{u}_N , contradicting that these distances are equal.

So $r_{-1} \neq r_1$. □

Claim 5.7. *If c is chaotic, then c is valence-unbalanced.*

Proof. We consider five cases.

Case 1. $V_0(c) \setminus \{(0, 0)\}$ is not contained in $y = x$, nor in one open half-plane of $y = -x$.

By Claim 5.4, there is a point \bar{v} of $V^*(c)$ on the line $y = -x$. There is a point $(k_0, -k_0) \in \text{conv}(V_0(c))$, $k_0 \neq 0$. Either $(0, 0)$ is between \bar{v} and $(k_0, -k_0)$ or it is between \bar{v} and $(-k_0, k_0)$; either way, it is in $V^*(c)$, so c is valence-unbalanced.

Case 2. $V_0(c) \setminus \{(0, 0)\}$ is not contained in $y = x$, but is contained in one open half-plane of $y = -x$.

There is $(k_0, k_1) \in V_0(c)$ such that $k_0 \neq k_1$ and $k_0 \neq -k_1$. By Claim 5.4, there is a point $\bar{v} \in V_1(c)$ strictly on the other side of $y = -x$. If \bar{v} is on $y = x$, then $(0, 0)$ is on the line segment between \bar{v} and $\bar{w} = (\frac{k_0+k_1}{2}, \frac{k_0+k_1}{2})$, hence in $V^*(c)$, so c is valence-unbalanced. If \bar{v} is not on $y = x$, then it has a reflection $\bar{u} \in V_{-1}(c)$. The set $\text{conv}\{\bar{v}, \bar{u}, \bar{w}\}$ contains an open disc about $(0, 0)$. Since there are points in $\text{conv}(V_0(c))$ arbitrarily close to \bar{w} on the line between (k_0, k_1) and (k_1, k_0) , we may choose (k'_0, k'_1) so $k'_0 \neq k'_1$ and $(0, 0) \in \text{conv}\{\bar{v}, \bar{u}, (k'_0, k'_1)\}$.

Suppose c is valence-balanced. Say \bar{v} is (v_0, v_1) . Then

$$(0, 0) = r\bar{v} + (1 - 2r)(k'_0, k'_1) + r\bar{u} = r(v_0 + v_1, v_1 + v_0) + (1 - 2r)(k'_0, k'_1),$$

so $k'_0 = \frac{-r(v_0+v_1)}{1-2r} = k'_1$, a contradiction. Therefore c is valence-unbalanced.

Case 3. $V_0(c)$ is contained in $y = x$, and $V_1(c)$ is contained in a line parallel to $y = x$.

Then c is partisan by Claim 5.3, contradicting that c is chaotic.

Case 4. $V_0(c)$ is contained in $y = x$, $\text{conv}(V_1(c))$ contains a line segment not parallel to $y = x$, and $V_1(c)$ has points on either side of $y = -x$.

Then there is a point of $y = -x$ strictly between two points of $\text{conv}(V_1(c))$ such that the line segment between them is not parallel to $y = x$. By Claim 5.6, $(0, 0) = r_{-1}\bar{v}_{-1} + r_1\bar{v}_1$ where $\bar{v}_\ell \in \text{conv}(V_\ell(c))$, $r_\ell \in [0, 1]_{\mathbb{Q}}$, $r_{-1} + r_1 = 1$, but $r_{-1} \neq r_1$, i.e. c is valence-unbalanced.

Case 5. $V_0(c)$ is contained in $y = x$, $\text{conv}(V_1(c))$ contains a line segment not parallel to $y = x$, and $V_1(c)$ is entirely on one side of $y = -x$.

Let \bar{v} be strictly between two points of $V_1(c)$ on a line segment not parallel to $y = x$. By Claim 5.6, the nearest point \bar{w} to \bar{v} on the line $y = x$ is $r_{-1}\bar{v}_{-1} + r_1\bar{v}_1$ where $\bar{v}_\ell \in \text{conv}(V_\ell(c))$, $r_\ell \in [0, 1]_{\mathbb{Q}}$, $r_{-1} + r_1 = 1$, but $r_{-1} \neq r_1$. By Claim 5.4, $V_0(c)$ has a point $\bar{v}_0 = (k, k)$ strictly on the other side of $y = -x$ from \bar{v} , and hence from \bar{w} . So for some $r \in [0, 1]_{\mathbb{Q}}$,

$$(0, 0) = r\bar{v}_0 + (1 - r)(r_{-1}\bar{v}_{-1} + r_1\bar{v}_1) = (1 - r)r_{-1}\bar{v}_{-1} + r\bar{v}_0 + (1 - r)r_1\bar{v}_1,$$

and $(1 - r)r_{-1} \neq (1 - r)r_1$, so c is valence-unbalanced. \square

Claim 5.8. *If $\mathcal{C} \subseteq \mathfrak{C}$ is symmetric and $c \in \mathcal{C}$ is valence-unbalanced, then for each $\{x, y\} \in \binom{X}{2}$ there is $d \in \text{maj-cl}(\mathcal{C})$ such that $\text{dom } d = \{\{x, y\}\}$.*

Proof. Suppose $\{x, y\} \in \binom{X}{2}$. There are $\bar{v}_\ell \in \text{conv}(V_\ell(c))$ and $r_\ell \in [0, 1]_{\mathbb{Q}}$, $r_{-1} + r_0 + r_1 = 1$, $r_{-1} \neq r_1$, so $r_{-1}\bar{v}_{-1} + r_0\bar{v}_0 + r_1\bar{v}_1 = (0, 0)$. Now,

$$\bar{v}_{-1} = \sum_{\bar{v} \in V_{-1}(c)} s_{\bar{v}}^{-1} \bar{v}, \quad \bar{v}_0 = \sum_{\bar{v} \in V_0(c)} s_{\bar{v}}^0 \bar{v}, \quad \bar{v}_1 = \sum_{\bar{v} \in V_1(c)} s_{\bar{v}}^1 \bar{v},$$

where $\sum_{\bar{v} \in V_\ell(c)} s_{\bar{v}}^\ell = 1$ for each ℓ . Each $\bar{v} \in V_\ell(c)$ is $(\text{val}_c(u_{\bar{v}}^\ell) - \ell, \text{val}_c(w_{\bar{v}}^\ell) + \ell)$ for some $\{u_{\bar{v}}^\ell, w_{\bar{v}}^\ell\} \in \binom{X}{2}$. Let $\sigma_{\bar{v}}^\ell$ be a permutation taking $u_{\bar{v}}^\ell$ to x and $w_{\bar{v}}^\ell$ to y , i.e.

$$\sigma_{\bar{v}}^\ell = \begin{cases} (x, y) & \text{if } u_{\bar{v}}^\ell = y, w_{\bar{v}}^\ell = x, \\ (x, w_{\bar{v}}^\ell, y) & \text{if } u_{\bar{v}}^\ell = y, w_{\bar{v}}^\ell \neq x, \\ (u_{\bar{v}}^\ell, x)(w_{\bar{v}}^\ell, y) & \text{otherwise.} \end{cases}$$

Let $\Gamma_{x,y}$ be all the permutations of X which fix x and y , and let

$$\bar{t} = \frac{1}{|\Gamma_{x,y}|} \sum_{\ell \in \{-1, 0, 1\}} r_\ell \sum_{\bar{v} \in V_\ell(c)} s_{\bar{v}}^\ell \sum_{\tau \in \Gamma_{x,y}} \text{pr}(c^{\tau \sigma_{\bar{v}}^\ell}).$$

Of course, the sum of the coefficients is

$$\sum_{\substack{\ell \in \{-1, 0, 1\} \\ \bar{v} \in V_\ell(c) \\ \tau \in \Gamma_{x,y}}} \frac{r_\ell s_{\bar{v}}^\ell}{|\Gamma_{x,y}|} = \frac{|\Gamma_{x,y}|}{|\Gamma_{x,y}|} \sum_{\ell \in \{-1, 0, 1\}} r_\ell = 1.$$

Since r_ℓ , $s_{\bar{v}}^\ell$, and $\frac{1}{|\Gamma_{x,y}|}$ are in $[0, 1]_{\mathbb{Q}}$, $\frac{r_\ell s_{\bar{v}}^\ell}{|\Gamma_{x,y}|} \in [0, 1]_{\mathbb{Q}}$. So $\bar{t} \in \text{pr-cl}(\mathcal{C})$.

Consider $t_{x,y}$. Each τ fixes x and y , so

$$\begin{aligned} c^{\tau\sigma_{\bar{v}}^\ell}\{x, y\} &= x \iff \\ c^{\sigma_{\bar{v}}^\ell}\{x, y\} &= x \iff \\ c\{u_{\bar{v}}^\ell, w_{\bar{v}}^\ell\} &= u_{\bar{v}}^\ell \iff \ell = 1. \end{aligned}$$

So $W_y^x(c^{\tau\sigma_{\bar{v}}^\ell}) = \ell$. Thus

$$\begin{aligned} t_{x,y} &= \frac{1}{|\Gamma_{x,y}|} \sum_{\ell \in \{-1, 0, 1\}} r_\ell \sum_{\bar{v} \in V_\ell(c)} s_{\bar{v}}^\ell \sum_{\tau \in \Gamma_{x,y}} \ell \\ &= \frac{|\Gamma_{x,y}|}{|\Gamma_{x,y}|} \sum_{\ell \in \{-1, 0, 1\}} \ell r_\ell = r_1 - r_{-1}. \end{aligned}$$

Since $r_{-1} \neq r_1$, $t_{x,y} \neq 0$.

Consider $t_{x,u}$ where $u \notin \{x, y\}$.

$$\begin{aligned} c^{\tau\sigma_{\bar{v}}^\ell}\{x, u\} &= x \iff \\ c^{\sigma_{\bar{v}}^\ell}\{x, \tau^{-1}(u)\} &= x \iff \\ c\{u_{\bar{v}}^\ell, u^*\} &= u_{\bar{v}}^\ell, \end{aligned}$$

where

$$u^* = (\sigma_{\bar{v}}^\ell)^{-1}(\tau^{-1}(u)) = \begin{cases} \tau^{-1}(u) & \text{if } \tau^{-1}(u) \notin \{u_{\bar{v}}^\ell, w_{\bar{v}}^\ell\}, \\ x & \text{if } \tau^{-1}(u) = u_{\bar{v}}^\ell, \\ y & \text{if } \tau^{-1}(u) = w_{\bar{v}}^\ell. \end{cases}$$

u^* is never $u_{\bar{v}}^\ell$ or $w_{\bar{v}}^\ell$. For a given ℓ and \bar{v} , u^* varies equally over all elements of X besides $\{u_{\bar{v}}^\ell, w_{\bar{v}}^\ell\}$, so

$$\begin{aligned} \sum_{\tau \in \Gamma_{x,y}} W_u^x(c^{\tau\sigma_{\bar{v}}^\ell}) &= \sum_{\tau \in \Gamma_{x,y}} W_{u^*}^{u_{\bar{v}}^\ell}(c) = \sum_{\tau \in \Gamma_{x,y,u}} \sum_{u^* \in X \setminus \{u_{\bar{v}}^\ell, w_{\bar{v}}^\ell\}} W_{u^*}^{u_{\bar{v}}^\ell}(c) \\ &= |\Gamma_{x,y,u}| \left(\text{val}_c(u_{\bar{v}}^\ell) - W_{u_{\bar{v}}^\ell}^{u_{\bar{v}}^\ell}(c) - W_{w_{\bar{v}}^\ell}^{u_{\bar{v}}^\ell}(c) \right) = |\Gamma_{x,y,u}| \left(\text{val}_c(u_{\bar{v}}^\ell) - \ell \right). \end{aligned}$$

This follows from the fact that, for each $u^* \in X \setminus \{u_{\bar{v}}^\ell, w_{\bar{v}}^\ell\}$, there are $|\Gamma_{x,y,u}|$ permutations τ which take u to u^* . For any u , $W_u^u(c) = 0$, and by choice of $u_{\bar{v}}^\ell$ and $w_{\bar{v}}^\ell$, $W_{w_{\bar{v}}^\ell}^{u_{\bar{v}}^\ell}(c) = \ell$.

Recall that the $u_{\bar{v}}^\ell$ were chosen so that $\text{val}_c(u_{\bar{v}}^\ell) - \ell$ is the first coordinate of \bar{v} . Recall also that the r_ℓ , $s_{\bar{v}}^\ell$, and valence pairs \bar{v} were chosen so

$$\sum_{\ell \in \{-1, 0, 1\}} r_\ell \sum_{\bar{v} \in V_\ell(c)} s_{\bar{v}}^\ell \bar{v} = (0, 0).$$

$$\begin{aligned}
t_{x,u} &= \frac{1}{|\Gamma_{x,y}|} \sum_{\ell \in \{-1,0,1\}} r_\ell \sum_{\bar{v} \in V_\ell(c)} s_{\bar{v}}^\ell \sum_{\tau \in \Gamma_{x,y}} W_u^x \left(c^{\tau \sigma_{\bar{v}}^\ell} \right) \\
&= \frac{1}{|\Gamma_{x,y}|} \sum_{\ell \in \{-1,0,1\}} r_\ell \sum_{\bar{v} \in V_\ell(c)} s_{\bar{v}}^\ell |\Gamma_{x,y,u}| (\text{val}_c(u_{\bar{v}}^\ell) - \ell) \\
&= 0.
\end{aligned}$$

Similarly, $t_{y,u} = 0$ for all $u \notin \{x, y\}$.

Consider $t_{u,w}$ with neither u nor w in $\{x, y\}$. Then

$$\begin{aligned}
c^{\tau \sigma_{\bar{v}}^\ell} \{u, w\} &= u \iff \\
c^{\sigma_{\bar{v}}^\ell} \{\tau^{-1}(u), \tau^{-1}(w)\} &= \tau^{-1}(u) \iff \\
c\{u^*, w^*\} &= u^*,
\end{aligned}$$

where $u^* = (\sigma_{\bar{v}}^\ell)^{-1}(\tau^{-1}(u))$ and $w^* = (\sigma_{\bar{v}}^\ell)^{-1}(\tau^{-1}(w))$. But $\tau^{-1}(u)$ and $\tau^{-1}(w)$ are never x or y , so u^* is never $u_{\bar{v}}^\ell$ and w^* is never $w_{\bar{v}}^\ell$. As τ varies, u^* and w^* vary equally over all members of X except $u_{\bar{v}}^\ell$ and $w_{\bar{v}}^\ell$, so $c\{u^*, w^*\} = u^*$ just as often as $c\{u^*, w^*\} = w^*$. Thus

$$\sum_{\tau \in \Gamma_{x,y}} W_w^u(c^{\tau \sigma_{\bar{v}}^\ell}) = 0.$$

Hence

$$t_{u,v} = \frac{1}{|\Gamma_{x,y}|} \sum_{\ell \in \{-1,0,1\}} r_\ell \sum_{\bar{v} \in V_\ell(c)} s_{\bar{v}}^\ell \sum_{\tau \in \Gamma_{x,y}} W_w^u(c^{\tau \sigma_{\bar{v}}^\ell}) = 0.$$

So $d = \text{maj}(\bar{t}) \in \text{maj-cl}(\mathcal{C})$ has $\text{dom } d = \{\{x, y\}\}$; all other pairs have $t_{u,v} = 0$. \square

Claim 5.9. *If $\mathcal{C} \subseteq \mathfrak{C}$ is symmetric and chaotic, then $\text{maj-cl}(\mathcal{C}) = \mathfrak{C}$.*

Proof. Suppose $f \in \mathfrak{C}$.

By Claim 5.7, there is some $c \in \mathcal{C}$ which is valence-unbalanced. So by Claim 5.8, for any $\{x, y\} \in \binom{X}{2}$ there is $d \in \text{maj-cl}(\mathcal{C})$ with $\text{dom } d = \{\{x, y\}\}$. If $d\{x, y\} = y$, by symmetry there is $d' \in \text{maj-cl}(\mathcal{C})$ with domain $\{\{x, y\}\}$ and $d'\{x, y\} = x$. So for every $\{x, y\} \in \text{dom } f$, let $d_{x,y} \in \text{maj-cl}(\mathcal{C})$ such that $\text{dom } d_{x,y} = \{\{x, y\}\}$ and $d_{x,y}\{x, y\} = f\{x, y\}$. Now combine the voter populations which yielded each $d_{x,y}$.

Since f was arbitrary, we have shown $\mathfrak{C} \subseteq \text{maj-cl}(\mathcal{C})$. Hence $\text{maj-cl}(\mathcal{C})$ is all of \mathfrak{C} . \square

6. Bounds

We shall consider the loose upper bounds, implied by the preceding proofs, on the necessary number of voters from a symmetric set to yield an arbitrary function in its majority closure, in terms of \mathbf{n} . There is no doubt that much tighter bounds could be obtained.

Trivial

0 functions suffice to obtain no outcome.

Balanced

In order to create a triangular function from a chosen function c , $(\mathbf{n} - 3)!$ permutations of the rest of the set were used. These were repeated 3 times, to establish each pair of edges in the triangle.

An arbitrary pseudo-balanced function has each edge in a directed cycle, so one cycle per edge suffices; so we have at most $\binom{\mathbf{n}}{2} = \frac{\mathbf{n}(\mathbf{n}-1)}{2}$ cycles. Each is constructed with two fewer triangles than the number of nodes it contains. This is at most $\mathbf{n} - 2$, if some cycle contains every node. So all together we have

$$\frac{1}{2}\mathbf{n}(\mathbf{n} - 1)(\mathbf{n} - 2)(\mathbf{n} - 3)! \cdot 3 = \frac{3\mathbf{n}!}{2}.$$

Partisan

There are at most \mathbf{n} tiers in an arbitrary tiered function, each with at most \mathbf{n} elements (of course, no function meets both conditions.) For each such element x , we take functions for each permutation holding x fixed; there are $(\mathbf{n} - 1)!$ such. So we have $\mathbf{n} \mathbf{n}(\mathbf{n} - 1)! = \mathbf{n} \mathbf{n}!$ as an upper bound.

Not balanced, not partisan, but not chaotic

The proof of Claim 4.5 gives the size of the population T_z yielding $c^{a,b,z}$ as $|T_z| = 3(\mathbf{n} - 3)!$, with $m = 2(\mathbf{n} - 3)!$ votes for the winner of each pair. We chose a partisan function c with l candidates in the winning tier. For each of at most $\binom{\mathbf{n}}{2}$ edges $a \rightarrow b$ and each of the $(\mathbf{n} - 2)$ $z \in X \setminus \{a, b\}$, we used k copies of T_z , where

$$k = \begin{cases} (\mathbf{n} - 2) \binom{\mathbf{n}-3}{l-1} & \text{if } l \leq \frac{\mathbf{n}}{2}, \\ (\mathbf{n} - 2) \binom{\mathbf{n}-3}{l-2} & \text{if } l > \frac{\mathbf{n}}{2}, \end{cases}$$

and m copies of a population of partisan functions D , where

$$|D| = \begin{cases} (\mathbf{n} - 2l) \binom{\mathbf{n}-1}{l} + (\mathbf{n} - 2) \binom{\mathbf{n}-2}{l-1} & \text{if } l \leq \frac{\mathbf{n}}{2}, \\ (2l - \mathbf{n}) \binom{\mathbf{n}-1}{l-1} + (\mathbf{n} - 2) \binom{\mathbf{n}-2}{l-1} & \text{if } l > \frac{\mathbf{n}}{2}. \end{cases}$$

In fact, $k' = \frac{k}{\gcd(m,k)}$ copies of T_z , and $m' = \frac{m}{\gcd(m,k)}$ copies of D , would suffice. Certainly $\binom{\mathbf{n}-3}{l-1} = \frac{(\mathbf{n}-3)!}{(l-1)!(\mathbf{n}-l-2)!} \mid (\mathbf{n} - 3)!$, and $\binom{\mathbf{n}-3}{l-2} = \frac{(\mathbf{n}-3)!}{(l-2)!(\mathbf{n}-l-1)!} \mid (\mathbf{n} - 3)!$, so we may take $k' = (\mathbf{n} - 2)$.

If $l \leq \frac{\mathbf{n}}{2}$, then we take $m' = 2(l - 1)!(\mathbf{n} - l - 2)!$, and

$$\begin{aligned} m' |D| &= 2(l - 1)!(\mathbf{n} - l - 2)! \left((\mathbf{n} - 2l) \binom{\mathbf{n} - 1}{l} + (\mathbf{n} - 2) \binom{\mathbf{n} - 2}{l - 1} \right) \\ &= 2(l - 1)!(\mathbf{n} - l - 2)! \left(\frac{(\mathbf{n} - 2l)(\mathbf{n} - 1)!}{l!(\mathbf{n} - l - 1)!} + \frac{(\mathbf{n} - 2)(\mathbf{n} - 2)!}{(l - 1)!(\mathbf{n} - l - 1)!} \right) \\ &= 2(\mathbf{n} - 2)! \left(\frac{(\mathbf{n} - 2l)(\mathbf{n} - 1)}{l(\mathbf{n} - l - 1)} + \frac{\mathbf{n} - 2}{\mathbf{n} - l - 1} \right). \end{aligned}$$

The last multiplicand is

$$\frac{\mathbf{n}(\mathbf{n} - l - 1)}{l(\mathbf{n} - l - 1)} = \frac{\mathbf{n}}{l} \leq \mathbf{n}.$$

If $l > \frac{\mathbf{n}}{2}$, then we take $m' = 2(l - 2)!(\mathbf{n} - l - 1)!$, and

$$\begin{aligned} m' |D| &= 2(l - 2)!(\mathbf{n} - l - 1)! \left((2l - \mathbf{n}) \binom{\mathbf{n} - 1}{l - 1} + (\mathbf{n} - 2) \binom{\mathbf{n} - 2}{l - 1} \right) \\ &= 2(l - 2)!(\mathbf{n} - l - 1)! \left(\frac{(2l - \mathbf{n})(\mathbf{n} - 1)!}{(l - 1)!(\mathbf{n} - l)!} + \frac{(\mathbf{n} - 2)(\mathbf{n} - 2)!}{(l - 1)!(\mathbf{n} - l - 1)!} \right) \\ &= 2(\mathbf{n} - 2)! \left(\frac{(2l - \mathbf{n})(\mathbf{n} - 1)}{(l - 1)(\mathbf{n} - l)} + \frac{\mathbf{n} - 2}{l - 1} \right). \end{aligned}$$

The last multiplicand is

$$\frac{\mathbf{n}(l - 1)}{(l - 1)(\mathbf{n} - l)} = \frac{\mathbf{n}}{\mathbf{n} - l} \leq \mathbf{n}.$$

So the total number of voters is

$$\begin{aligned} &\binom{\mathbf{n}}{2} ((\mathbf{n} - 2) |T_z| k' + m' |D|) \\ &\leq \frac{\mathbf{n}(\mathbf{n} - 1)}{2} ((\mathbf{n} - 2)3(\mathbf{n} - 3)!(\mathbf{n} - 2) + 2(\mathbf{n} - 2)!\mathbf{n}) \\ &= \mathbf{n}(\mathbf{n} - 1)(\mathbf{n} - 2)! \left(\frac{3}{2}(\mathbf{n} - 2) + \mathbf{n} \right) \\ &= \mathbf{n}! \left(\frac{3}{2}\mathbf{n} - 3 + \mathbf{n} \right) < \frac{5}{2}\mathbf{n}!. \end{aligned}$$

Chaotic

Given an arbitrary function, we wish to create a “single-edged” $d_{x,y}$, as developed in Section 5, for each edge in the function; there are at most $\binom{\mathbf{n}}{2}$ of these. In creating such a single-edged $d_{x,y}$, we use a collection of permutations of a choice function, under permutations which take x and y to elements having certain valence combinations. Reading the proof of Claim 5.7 carefully, we see that at most 4 such valence pairs are used. For each of these, all permutations fixing the relevant pair are used, giving $4(\mathbf{n} - 2)!$ choice functions used in the creation of the single-edged function. However, these are being combined by some set of rational coefficients, so in fact we must expand each such function taking as many copies as the numerator of its associated coefficient when put over the least common denominator L . The coefficients sum to 1, so the sum of the numbers of copies is L ; so $L(\mathbf{n} - 2)!$ functions are used for each edge.

Six possible linear combinations of valence pairs yielding $(0, 0)$ were considered in Claim 5.7. Using the fact that the original valence pairs are pairs of integers between $-\mathbf{n}$ and \mathbf{n} , we can solve for the coefficients by matrix inversion

and determine that the least common denominator is at most the determinant of the associated matrix. For instance, in Case 4, suppose there is a point of $y = -x$ strictly between two points of $V_1(c)$ on a line segment not parallel to $y = x$. (This is true in this Case, unless $V_1(c)$ is contained in a line parallel to $y = x$ except for a point on the line $y = -x$; such a situation actually has a smaller common denominator.)

We use Claim 5.6, with $(0,0)$ for \bar{w} , and note that either the line from \bar{p}_0 through \bar{w} meets the line L , or the line from \bar{p}_1 through \bar{w} does. Say \bar{p} is the one which works. Then $(0,0)$ is an unbalanced linear combination of the three valence pairs \bar{p} , \bar{q}_0 , and \bar{q}_1 . Say $\bar{p} = (a, b)$; then one of the \bar{q} is (b, a) and the other is (c, d) . Solving

$$\begin{aligned} r + s + t &= 1 \\ ar + bs + ct &= 0 \\ br + as + dt &= 0 \end{aligned}$$

amounts to

$$\begin{bmatrix} r \\ s \\ t \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ a & b & c \\ b & a & d \end{bmatrix}^{-1} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}.$$

Since everything on the right hand side is an integer, the only denominator introduced is the determinant of the 3 by 3 matrix, $bd - ac - ad + bc + a^2 - b^2$. Since each of a, b, c, d is an integer between $-\mathbf{n}$ and \mathbf{n} , this is at most $6\mathbf{n}^2$.

In the cases involving four points, such as Case 1 or Case 5, an additional constraint is needed, which is not a consequence of the other three. In Case 1, use $r_1 + r_2 + r_3 + r_4 = 1$, $a_1r_1 + a_2r_2 + a_3r_3 + a_4r_4 = 0$, $b_1r_1 + b_2r_2 + b_3r_3 + b_4r_4 = 0$, and add $b_3r_3 + b_4r_4 = -a_3r_3 - a_4r_4$, since the points \bar{v} and $(k_0, -k_0)$ were linear combinations of two valence pairs on the line $y = -x$. The lowest common denominator is at most the determinant of

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \\ 0 & 0 & a_3 + b_3 & a_4 + b_4 \end{bmatrix},$$

which expands to a 16-term sum of three a_i or b_i each, hence at most $16\mathbf{n}^3$. In Case 5, note that the point \bar{v} between two points of $V_1(c)$ is arbitrary; we can choose any point between the two. Between any two pairs of integers on which do not lie on a line perpendicular to $y = x$, we can choose a point whose projection on $y = x$ is of the form $(\frac{l}{4}, \frac{l}{4})$ for some integer l . Solving

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ k & a & b & c \\ k & b & a & d \\ 0 & a & b & c \end{bmatrix}^{-1} \begin{bmatrix} 1 \\ 0 \\ 0 \\ \frac{l}{4} \end{bmatrix}$$

gives a determinant $k(2a^2 - b^2 - ab - ac + bd + bc - ad)$ at most $8\mathbf{n}^3$; we must multiply by 4 because of the fraction in the column vector, giving $32\mathbf{n}^3$. This is the largest denominator of the six possible linear combinations in Claim 5.7.

Thus our upper bound is

$$L(\mathbf{n} - 2)! \frac{\mathbf{n}^2 - \mathbf{n}}{2} < 16 \mathbf{n}^3 \mathbf{n}(\mathbf{n} - 1)(\mathbf{n} - 2)! = 16 \mathbf{n}^3 \mathbf{n}!$$

- [1] S. Shelah, What majority decisions are possible, *Discrete Mathematics* 309 (2009) 2349–2364.
- [2] J. A. N. d. Condorcet, *Essai sur l'application de l'analyse à la probabilité des décisions rendues à la pluralité des voix*, L'imprimerie royale, Paris, 1785.
- [3] D. C. McGarvey, A theorem on the construction of voting paradoxes, *Econometrica* 21 (1953) 608–610.
- [4] R. Stearns, The voting problem, *American Mathematical Monthly* 66 (1959) 761–763.
- [5] P. Erdős, L. Moser, On the representation of directed graphs as unions of orderings, *Magyar Tud. Akad. Mat. Kutats Int. Kvzl.* 9 (1964) 125–132.
- [6] J. Mala, On λ -majority voting paradoxes, *Mathematical Social Sciences* 37 (1999) 39–44.
- [7] N. Alon, Voting paradoxes and digraphs realizations, *Advances in Appl. Math.* 29 (2002) 126–135.
- [8] K.-I. Inada, A note on the simple majority decision rule, *Econometrica* 32 (1964) 525–553.